



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
28.07.1999 Bulletin 1999/30

(51) Int Cl.⁶: **G01N 33/576, C07K 14/18**

(21) Application number: **95114016.9**

(22) Date of filing: **03.04.1991**

(54) **Combinations of hepatitis C virus (HCV) antigens for use in immunoassays for anti-HCV antibodies**

Kombinationen der Hepatitis-C-Virus (HCV) Antigene zur Verwendung in Immuntests für Anti-HCV-Antikörper

Combinaisons d'antigènes de virus hépatite-C (HCV) pour l'utilisation dans des essais immunologiques d'anticorps d'anti-HCV

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(30) Priority: **04.04.1990 US 504352**

(43) Date of publication of application:
24.01.1996 Bulletin 1996/04

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
91302910.4 / 0 450 931

(73) Proprietor: **CHIRON CORPORATION**
Emeryville, California 94608 (US)

(72) Inventors:

- **Houghton, Michael**
Danville, California 94526 (US)
- **Choo, Qui-Lim**
El Cerrito, California 94530 (US)
- **Kuo, George**
San Francisco, California 94112 (US)

(74) Representative: **Goldin, Douglas Michael**
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5LX (GB)

(56) References cited:
EP-A- 0 318 216 **WO-A-89/04669**

EP 0 693 687 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionTechnical Field

5 **[0001]** The present invention is in the field of immunoassays for HCV (previously called Non-A, Non-B hepatitis virus). More particularly, it concerns combinations of HCV antigens that permit broad range immunoassays for anti-HCV antibodies.

Background

10 **[0002]** The disease known previously as Non-A, Non-B hepatitis (NANBH) was considered to be a transmissible disease or family of diseases that were believed to be viral-induced, and that were distinguishable from other forms of viral-associated liver diseases, including that caused by the known hepatitis viruses, i.e., hepatitis A virus (HAV), hepatitis B virus (HBV), and delta hepatitis virus (HDV), as well as the hepatitis induced by cytomegalovirus (CMV) or Epstein-Barr virus (EBV). NANBH was first identified in transfused individuals. Transmission from man to chimpanzee and serial passage in chimpanzees provided evidence that NANBH was due to a transmissible infectious agent or agents. Epidemiologic evidence suggested that there may be three types of NANBH: a water-borne epidemic type; a blood-borne or parenterally transmitted type; and a sporadically occurring (community acquired) type. However, until recently, no transmissible agent responsible for NANBH had been identified, and clinical diagnosis and identification of NANBH had been accomplished primarily by exclusion of other viral markers. Among the methods used to detect putative NANBH antigens and antibodies were agar-gel diffusion, counterimmunoelectrophoresis, immunofluorescence microscopy, immune electron microscopy, radioimmunoassay, and enzyme-linked immunosorbent assay. However, none of these assays proved to be sufficiently sensitive, specific, and reproducible to be used as a diagnostic test for NANBH.

25 **[0003]** In 1987, scientists at Chiron Corporation (the owner of the present application) identified the first nucleic acid definitively linked to blood-borne NANBH. See, e.g., EPO Pub. No. 318,216; Houghton et al., Science 244: 359 (1989). These publications describe therein the cloning of an isolate from a new viral class being named hepatitis C virus (HCV) the prototype virus described therein being named "HCV1." HCV is a flavi-like virus, with an RNA genome. EP-A-318,216 also describes the immunogenicity of the C100 polypeptide expressed from HCV cDNA and the use of HCV antigens in diagnostic immunoassays.

30 **[0004]** U.S. Patent No. 5,350,671 (Houghton et al.), describes the preparation of various recombinant HCV polypeptides by expressing HCV cDNA and the screening of those polypeptides for immunological reactivity with sera from HCV patients. That limited screening showed that at least five of the polypeptides tested were very immunogenic; specifically, those identified as 5-1-1, C100, C33c, CA279c, and CA290a. Of these five polypeptides, 5-1-1 is located in the putative NS4 domain; C100 spans the putative NS3 and NS4 domains; C33c is located within the putative NS3 domain and CA279a and CA290a are located within the putative C domain. The screening also showed that no single polypeptide tested was immunologically reactive with all sera. Thus, improved tests, which react with all or more samples from HCV positive individuals are desirable. WO 91/15574, published 17th October 1991, describes purified proteins and glycoproteins of HCV useful in a diagnostic system for detection of human HCV antisera.

Disclosure of the Invention

45 **[0005]** Applicants have carried out additional serological studies on HCV antigens that confirm that no single HCV polypeptide identified to date is immunologically reactive with all sera. This lack of a single polypeptide that is universally reactive with all sera from individuals with HCV may be due, inter alia, to strain-to-strain variation in HCV epitopes, variability in the humoral response from individual-to-individual and/or variation in serology with the state of the disease.

[0006] These additional studies have also enabled applicants to identify combinations of HCV antigens that provide more efficient detection of HCV antibodies than any single HCV polypeptide.

50 **[0007]** Accordingly, one aspect of this invention is a combination of hepatitis C virus (HCV) antigens in one or more polypeptides made by chemical synthesis or recombinant expression, comprising:

- (a) a first HCV antigen comprising an epitope from the C domain of the HCV polyprotein; and
- (b) a second HCV antigen selected from:

- 55 (i) an HCV antigen comprising an epitope from the S domain;
- (ii) an HCV antigen comprising an epitope from the NS3 domain; (iii) an HCV antigen comprising an epitope from the NS4 domain; and
- (iv) an HCV antigen comprising an epitope from the NS5 domain;

with the proviso that the combination is not the peptide p1 with C100-3, the peptide p35 with C100-3, the peptide p99 with C100-3 or a peptide having amino acids 9 to 177 of the HCV-1 polyprotein.

[0008] Preferably the second HCV antigen comprises an epitope from the S domain and in this case, the combination preferably further comprises at least one additional HCV antigen selected from:

- (i) an HCV antigen comprising an epitope from the NS3 domain;
- (ii) an HCV antigen comprising an epitope from the NS4 domain; and
- (iii) an HCV antigen comprising an epitope from the NS5 domain;

More preferably, the additional antigen is from the NS3 domain.

[0009] Another aspect of the invention is a method for detecting antibodies to HCV in a mammalian body component suspected of containing said antibodies comprising contacting said body component with the above-described combination of HCV antigens under conditions that permit antibody-antigen reaction and detecting the presence of immune complexes of said antibodies and said antigens.

[0010] Another aspect of the invention is a method for detecting antibodies to HCV in a mammalian body component suspected of containing said antibodies comprising contacting said body component with a combination of HCV antigens of the invention under conditions that permit antibody-antigen reaction and detecting the presence of immune complexes of said antibodies and said antigens.

[0011] Another aspect of the invention is a kit for carrying out an assay for detecting antibodies to HCV in a mammalian body component suspected of containing said antibodies comprising in packaged combination.

- (a) said combination of HCV antigens
- (b) standard control reagents; and
- (c) instructions for carrying out the assay.

Brief Description of the Drawings

[0012] In the drawings:

[0013] Figure 1 is the nucleotide sequence of the cDNA sense and anti-sense strand for the HCV polyprotein and the amino acid sequence encoded by the sense strand.

[0014] Figure 2 is a schematic of the amino acid sequence of Figure 1 showing the putative domains of the HCV polypeptide.

Modes for Carrying Out the Invention

Definitions

[0015] "HCV antigen" means a polypeptide of at least about 5 amino acids, more usually at least about 8 to 10 amino acids that defines an epitope found in an isolate of HCV. Preferably, the epitope is unique to HCV. When an antigen is designated by an alphanumeric code, the epitope is from the HCV domain specified by the alphanumeric.

[0016] "Synthetic" as used to characterize an HCV antigen means that the HCV antigen has been man-made such as by chemical or recombinant synthesis.

[0017] "Domains" means those segments of the HCV polyprotein shown in Figure 2 which generally correspond to the putative structural and nonstructural proteins of HCV. Domain designations generally follow the convention used to name Flaviviral proteins. The locations of the domains shown in Figure 2 are only approximate. The designations "NS" denotes "nonstructural" domains, while "S" denotes the envelope domain, and "C" denotes the nucleocapsid or core domain.

[0018] "Fusion polypeptide" means a polypeptide in which the HCV antigen(s) are part of a single continuous chain of amino acids, which chain does not occur in nature. The HCV antigens may be connected directly to each other by peptide bonds or be separated by intervening amino acid sequences. The fusion polypeptides may also contain amino acid sequences exogenous to HCV.

[0019] "Common solid matrix" means a solid body to which the individual HCV antigens or the fusion polypeptide comprised of HCV antigens are bound covalently or by noncovalent means such as hydrophobic adsorption.

[0020] "Mammalian body component" means a fluid or tissue of a mammalian individual (e.g., a human) that commonly contains antibodies produced by the individual. Such components are known in the art and include, without limitation, blood, plasma, serum, spinal fluid, lymph fluid, secretions of the respiratory, intestinal or genitourinary tracts, tears, saliva, milk, white blood cells, and myelomas.

[0021] "Immunologically reactive" means that the antigen in question will react specifically with anti-HCV antibody commonly present in a significant proportion of sera from individuals infected with HCV.

[0022] "Immune complex" means the combination or aggregate formed when an antibody binds to an epitope on an antigen.

Combinations of HCV Antigens

[0023] Figure 2 shows the putative domains of the HCV polyprotein. The domains from which the antigens used in the combinations derive are: C, S (or E), NS3, NS4, and NS5. The C domain is believed to define the nucleocapsid protein of HCV. It extends from the N-terminal of the polyprotein to approximately amino acid 120 of Figure 1. The S domain is believed to define the virion envelope protein, and possibly the matrix (M) protein, and is believed to extend from approximately amino acid 120 to amino acid 400 of Figure 1. The NS3 domain extends from approximately amino acid 1050 to amino acid 1640 and is believed to constitute the viral protease. The NS4 domain extends from the terminus of NS3 to approximately amino acid 2000. The function of the NS4 protein is not known at this time. Finally, the NS5 domain extends from about amino acid 2000 to the end of the polyprotein and is believed to define the viral polymerase.

[0024] The sequence shown in Figure 1 is the sequence of the HCV1 isolate. It is expected that the sequences of other strains of the blood-borne HCV may differ from the sequence of Figure 1, particularly in the envelope (S) and nucleocapsid (C) domains. The use of HCV antigens having such differing sequences is intended to be within the scope of the present invention, provided, however, that the variation does not significantly degrade the immunological reactivity of the antigen to sera from persons infected with HCV.

[0025] In general, the HCV antigens will comprise entire or truncated domains, the domain fragments being readily screened for antigenicity by those skilled in the art. The individual HCV antigens used in the combination will preferably comprise the immunodominant portion (i.e., the portion primarily responsible for the immunological reactivity of the polypeptide) of the stated domain. In the case of the C domain it is preferred that the C domain antigen comprise a majority of the entire sequence of the domain. The antigen designated C22 (see Example 4, *infra*), is particularly preferred. The S domain antigen preferably includes the hydrophobic subdomain at the N-terminal end of the domain. This hydrophobic subdomain extends from approximately amino acid 199 to amino acid 328 of Figure 1. The HCV antigen designated S2 (see Example 3, *infra*), is particularly preferred. Sequence downstream of the hydrophobic subdomain may be included in the S domain antigen if desired.

[0026] A preferred NS3 domain antigen is the antigen designated C33c. That antigen includes amino acids 1192 to 1457 of Figure 1. A preferred NS4 antigen is C100 which comprises amino acids 1569 to 1931 of Figure 1. A preferred NS5 antigen comprises amino acids 2054 to 2464 of Figure 1.

[0027] The HCV antigen may be in the form of a polypeptide composed entirely of HCV amino acid sequence or it may contain sequence exogenous to HCV (i.e., it may be in the form of a fusion protein that includes exogenous sequence). In the case of recombinantly produced HCV antigen, producing the antigen as a fusion protein such as with SOD, alpha-factor or ubiquitin (see commonly owned U.S. Pat. No. 4,751,180, U.S. Pat. No. 4,870,008 and U.S. Pat. Application Serial. No. 390,599, filed 7 August 1989, which describe expression of SOD, alpha-factor and ubiquitin fusion proteins) may increase the level of expression and/or increase the water solubility of the antigen. Fusion proteins such as the alpha-factor and ubiquitin fusion are processed by the expression host to remove the heterologous sequence. Alpha-factor is a secretion system, however, while ubiquitin fusions remain in the cytoplasm.

[0028] Further, the combination of antigens may be produced as a fusion protein. For instance, a continuous fragment of DNA encoding C22 and C33c may be constructed, cloned into an expression vector and used to express a fusion protein of C22 and C33c. In a similar manner fusion proteins of C22 and C100; C22 and S2; C22 and an NS5 antigen; C22, C33c, and S2; C22, C100 and S2, and C22, C33c, C100, and S2 may be made. Alternative fragments from the exemplified domain may also be used.

Preparation of HCV Antigens

[0029] The HCV antigens of the invention are produced recombinantly or by known solid phase chemical synthesis.

[0030] When produced by recombinant techniques, standard procedures for constructing DNA encoding the antigen, cloning that DNA into expression vectors, transforming host cells such as bacteria, yeast, insect, or mammalian cells, and expressing such DNA to produce the antigen may be employed. As indicated previously, it may be desirable to express the antigen as a fusion protein to enhance expression, facilitate purification, or enhance solubility. Examples of specific procedures for producing representative HCV antigens are described in the Examples, *infra*, and in US Patent 5,350,671.

Formulation of Antigens for Use in Immunoassay

[0031] The HCV antigens may be combined by producing them in the form of a fusion protein composed of two or more of the antigens, by immobilizing them individually on a common solid matrix, or by physically mixing them. Fusion proteins of the antigen may also be immobilized on (bound to) a solid matrix. Methods and means for covalently or noncovalently binding proteins to solid matrices are known in the art. The nature of the solid surface will vary depending upon the assay format. For assays carried out in microtiter wells, the solid surface will be the wall of the well or cup. For assays using beads, the solid surface will be the surface of the bead. In assays using a dipstick (i.e., a solid body made from a porous or fibrous material such as fabric or paper) the surface will be the surface of the material from which the dipstick is made. In agglutination assays the solid surface may be the surface of latex or gelatin particles. When individual antigens are bound to the matrix they may be distributed homogeneously on the surface or distributed thereon in a pattern, such as bands so that a pattern of antigen binding may be discerned.

[0032] Simple mixtures of the antigens comprise the antigens in any suitable solvent or dispersing medium.

Assay Formats Using Combinations of Antigens

[0033] The HCV antigens may be employed in virtually any assay format that employs a known antigen to detect antibodies. A common feature of all of these assays is that the antigen is contacted with the body component suspected of containing HCV antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

[0034] Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

[0035] The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type. In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidene fluoride (known as Immulon™), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immulon™ 1 or Immulon™ 2 microtiter plates or 6.4 mm (0.25 inch) polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.

[0036] In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

[0037] In a standard format, the amount of HCV antibodies forming the antibody-antigen complex is directly monitored. This may be accomplished by determining whether labeled anti-xenogenic (e.g., anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

[0038] Complexes formed comprising anti-HCV antibody (or, in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of antixenogenic Ig complexed with a label, (e.g., an enzyme label).

[0039] In an immunoprecipitation or agglutination assay format the reaction between the HCV antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-HCV antibody is present in the test specimen, no visible precipitate is formed.

[0040] The HCV antigens will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the combination of antigens (either already bound to a solid matrix or separate with reagents for binding them to the matrix), control antibody formulations (positive and/or negative), labeled antibody when the assay format requires same and signal generating reagents (e.g., enzyme substrate) if the label does not generate a signal directly. Instructions (e.g., written, tape, VCR, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

[0041] The following examples are intended to illustrate the invention and are not intended to limit the invention in any manner.

Example 1: Synthesis of HCV Antigen C33c

[0042] HCV antigen C33c contains a sequence from the NS3 domain. Specifically, it includes amino acids 1192-1457 of Figure 1. This antigen was produced in bacteria as a fusion protein with human superoxide dismutase (SOD) as follows. The vector pSODc1 (Steiner et al. (1986), J. Virol. 58:9) was digested to completion with EcoRI and BamHI and the resulting EcoRI, BamHI fragment was ligated to the following linker to form pcf1EF:

GATC CTG GAA TTC TGA TAA
GAC CTT AAG ACT ATT TTA A

A cDNA clone encoding amino acids 1192-1457 and having EcoRI ends was inserted into pcf1EF to form pcf1EF/C33c. This expression construct was transformed into D1210 *E. coli* cells.

[0043] The transformants were used to express a fusion protein comprised of SOD at the N-terminus and in-frame C33c HCV antigen at the C-terminus. Expression was accomplished by inoculating 1500 ml of Luria broth containing ampicillin (100 micrograms/ml) with 15 ml of an overnight culture of the transformants. The cells were grown to an O.D. of 0.3, IPTG was added to yield a final concentration of 2 mM, and growth continued until the cells attained a density of 1 O.D., at which time they were harvested by centrifugation at 3,000 x g at 4°C for 20 minutes. The packed cells can be stored at -80°C for several months.

[0044] In order to purify the SOD-C33c polypeptide the bacterial cells in which the polypeptide was expressed were subjected to osmotic shock and mechanical disruption, the insoluble fraction containing SOD-C33c was isolated and subjected to differential extraction with an alkaline-NaCl solution, and the fusion polypeptide in the extract purified by chromatography on columns of S-SepharoseTM and Q-SepharoseTM.

[0045] The crude extract resulting from osmotic shock and mechanical disruption was prepared by the following procedure. One gram of the packed cells were suspended in 10 ml of a solution containing 0.02 M Tris HCl, pH 7.5, 10 mM EDTA, 20% sucrose, and incubated for 10 minutes on ice. The cells were then pelleted by centrifugation at 4,000 x g for 15 min at 4°C. After the supernatant was removed, the cell pellets were resuspended in 10 ml of Buffer A1 (0.01M Tris HCl, pH 7.5, 1 mM EDTA, 14 mM betamercaptoethanol [BME]), and incubated on ice for 10 minutes. The cells were again pelleted at 4,000 x g for 15 minutes at 4°C. After removal of the clear supernatant (periplasmic fraction I), the cell pellets were resuspended in Buffer A1, incubated on ice for 10 minutes, and again centrifuged at 4,000 x g for 15 minutes at 4°C. The clear supernatant (periplasmic fraction II) was removed, and the cell pellet resuspended in 5 ml of Buffer A2 (0.02 M Tris HCl, pH 7.5, 14 mM BME, 1 mM EDTA, 1 mM PMSF). In order to disrupt the cells, the suspension (5 ml) and 7.5 ml of Dyno-millTM lead-free acid washed glass beads (0.10-0.15 mm diameter) (obtained from Glen-Mills, Inc.) were placed in a FalconTM tube, and vortexed at top speed for two minutes, followed by cooling for at least 2 min on ice; the vortexing-cooling procedure was repeated another four times. After vortexing, the slurry was filtered through a scintered glass funnel using low suction; the glass beads were washed two times with Buffer A2, and the filtrate and washes combined.

[0046] The insoluble fraction of the crude extract was collected by centrifugation at 20,000 x g for 15 min at 4°C, washed twice with 10 ml Buffer A2, and resuspended in 5 ml of MILLI-QTM water.

[0047] A fraction containing SOD-C33c was isolated from the insoluble material by adding to the suspension NaOH (2 M) and NaCl (2 M) to yield a final concentration of 20 mM each, vortexing the mixture for 1 minute, centrifuging it 20,000 x g for 20 min at 4°C, and retaining the supernatant.

[0048] In order to purify SOD-C33c on S-SepharoseTM, the supernatant fraction was adjusted to a final concentration of 6M urea, 0.05M Tris HCl, pH 7.5, 14 mM BME, 1 mM EDTA. This fraction was then applied to a column of S-SepharoseTM Fast Flow (1.5 x 10 cm) which had been equilibrated with Buffer B (0.05M Tris HCl, pH 7.5, 14 mM BME, 1 mM EDTA). After application, the column was washed with two column volumes of Buffer B. The flow through and wash fractions were collected. The flow rate of application and wash, was 1 ml/min; and collected fractions were 1 ml. In order to identify fractions containing SOD-C33c, aliquots of the fractions were analyzed by electrophoresis on 10% polyacrylamide gels containing SDS followed by staining with Coomassie blue. The fractions are also analyzable by Western blots using an antibody directed against SOD. Fractions containing SOD-C33c were pooled.

[0049] Further purification of SOD-C33c was on a Q-SepharoseTM column (1.5 x 5 cm) which was equilibrated with Buffer B. The pooled fractions containing SOD-C33c obtained from chromatography on S-Sepharose was applied to the column. The column was then washed with Buffer B, and eluted with 60 ml of a gradient of 0.0 to 0.4 M NaCl in Buffer B. The flow rate for application, wash, and elution was 1 ml/min; collected fractions were 1 ml. All fractions from the Q-SepharoseTM column were analyzed as described for the S-SepharoseTM column. The peak of SOD-C33c eluted from the column at about 0.2 M NaCl.

[0050] The SOD-C33c obtained from the Q-SepharoseTM column was greater than about 90% pure, as judged by

analysis on the polyacrylamide SDS gels and immunoblot using a monoclonal antibody directed against human SOD.

Example 2: Synthesis of HCV Antigen C100

5 [0051] HCV antigen C100 contains sequences from the NS3 and NS4 domains. Specifically, it includes amino acids 1569-1931 of Figure 1. This antigen was produced in yeast. A cDNA fragment of a 1270 bp encoding the above amino acids and heaving EcoRI termini was prepared.

10 [0052] The construction of a yeast expression vector in which this fragment was fused directly to the *S. cerevisiae* ADH2/GAP promoter was accomplished by a protocol which included amplification of the C100 sequence using a PCR method, followed by ligation of the amplified sequence into a cloning vector. After cloning, the C100 sequence was excised, and with a sequence which contained the ADH2/GAP promoter, was ligated to a large fragment of a yeast vector to yield a yeast expression vector.

15 [0053] The PCR amplification of C100 was performed using as template the vector pS3-56_{C100m}, which had been linearized by digestion with Sall. pS3-56, which is a pBR322 derivative, contains an expression cassette which is comprised of the ADH2/GAPDH hybrid yeast promoter upstream of the human superoxide dismutase gene, and a downstream alpha factor transcription terminator.

20 [0054] The oligonucleotide primers used for the amplification were designed to facilitate cloning into the expression vector, and to introduce a translation termination codon. Specifically, novel 5'-HindIII and 3'-Sall sites were generated with the PCR oligonucleotides. The oligonucleotide containing the Sall site also encodes the double termination codons, TAA and TGA. The oligonucleotide containing the HindIII site also contains an untranslated leader sequence derived from the pgap63 gene, situated immediately upstream of the AUG codon. The pEco63GAPDH gene is described by Holland and Holland (1980) and by Kniskern et al. (1986). The PCR primer sequences used for the direct expression of C100m were:

25
5' GAG TGC TCA AGC TTC AAA ACA AAA TGG CTC
ACT TTC TAT CCC AGA CAA AGC AGA GT 3'

30 and

35
5' GAG TGC TCG TCG ACT CAT TAG GGG GAA
ACA TGG TTC CCC CGG GAG GCG AA 3'.

40 [0055] Amplification by PCR, utilizing the primers, and template, was with a Cetus-Perkin-Elmer PCR kit, and was performed according to the manufacturer's directions. The PCR conditions were 29 cycles of 94°C for a minute, 37°C for 2 minutes, 72°C for 3 minutes; and the final incubation was at 72°C for 10 minutes. The DNA can be stored at 4°C or -20°C overnight.

45 [0056] After amplification, the PCR products were digested with HindIII and Sall. The major product of 1.1 kb was purified by electrophoresis on a gel, and the eluted purified product was ligated with a large Sall-HindIII fragment of pBR322. In order to isolate correct recombinants, competent HB101 cells were transformed with the recombinant vectors, and after cloning, the desired recombinants were identified on the basis of the predicted size of HindIII-Sall fragments excised from the clones. One of the clones which contained the a HindIII-Sall fragment of the correct size was named pBR322/C100-d. Confirmation that this clone contained amplified C100 was by direct sequence analysis of the HindIII-Sall fragment.

50 [0057] The expression vector containing C100 was constructed by ligating the HindIII-Sall fragment from pBR322/C100-d to a 13.1 kb BamHI-Sall fragment of pBS24.1, and a 1369 bp BamHI-HindIII fragment containing the ADH2/GAP promoter. (The latter fragment is described in EPO 164,556). The ADH2/GAP promoter fragment was obtained by digestion of the vector pPGAP/AG/HindIII with HindIII and BamHI, followed by purification of the 1369 bp fragment on a gel.

55 [0058] Competent HB101 cells were transformed with the recombinant vectors; and correct recombinants were identified by the generation of a 2464 bp fragment and a 13.1 kb fragment generated by BamHI and Sall digestion of the cloned vectors. One of the cloned correct recombinant vectors was named pC100-d#3.

[0059] In order to express C100, competent cells of *Saccharomyces cerevisiae* strain AB122 (MATa leu2 ura3-53 prb 1-1122 pep4-3 prcl-407[cir-0]) were transformed with the expression vector pC100-d#3. The transformed cells were plated on URA-sorbitol, and individual transformants were then streaked on Leu⁻ plates.

[0060] Individual clones were cultured in Leu^r, ura^r medium with 2% glucose at 30°C for 24-36 hours. One liter of Yeast Extract Peptone Medium (YEP) containing 2% glucose was inoculated with 10 ml of the overnight culture, and the resulting culture was grown at 30°C at an agitation rate of 400 rpm and an aeration rate of 1 L of air per 1 L of medium per minute (i.e., 1vvm) for 48 hours. The pH of the medium was not controlled. The culture was grown in a BioFlo II fermentor manufactured by New Brunswick Science Corp. Following fermentation, the cells were isolated and analyzed for C100 expression.

[0061] Analysis for expressed C100 polypeptide by the transformed cells was performed on total cell lysates and crude extracts prepared from single yeast colonies obtained from the Leu^r plates. The cell lysates and crude extracts were analyzed by electrophoresis on SDS polyacrylamide gels, and by Western blots. The Western blots were probed with rabbit polyclonal antibodies directed against the SOD-C100 polypeptide expressed in yeast. The expected size of the C100 polypeptide is 364 amino acids. By gel analysis the expressed polypeptide has a MW_r of 39.9K.

[0062] Both analytical methods demonstrated that the expressed C100 polypeptide was present in total cell lysates, but was absent from crude extracts. These results suggest that the expressed C100 polypeptide may be insoluble.

Example 3: Expression of HCV Antigen S2

[0063] HCV antigen S2 contains a sequence from the hydrophobic N-terminus of the S domain. It includes amino acids 199-328 of Figure 1.

[0064] The protocol for the construction of the expression vector encoding the S2 polypeptide and for its expression in yeast was analogous to that used for the expression of the C100 polypeptide, described in Example 2.

[0065] The template for the PCR reaction was the vector pBR322/Pi14a, which had been linearized by digestion with HindIII. Pi14a is a cDNA clone that encodes amino acids 199-328.

[0066] The oligonucleotides used as primers for the amplification by PCR of the S2 encoding sequence were the following.

For the 5'-region of the S2 sequence:

5' GAG TGC TCA AGC TTC AAA ACA AAA TGG GGC TCT
ACC ACG TCA CCA ATG ATT GCC CTA AC 3' ;

and

for the 3'-region of the S2 sequence:

5' GAG TGC TCG TCG ACT CAT TAA GGG GAC CAG TTC
ATC ATC ATA TCC CAT GCC AT 3' .

The primer for the 5'-region introduces a HindIII site and an ATG start codon into the amplified product. The primer for the 3'-region introduces translation stop codons and a Sall site into the amplified product.

[0067] The PCR conditions were 29 cycles of 94°C for a minute, 37°C for 2 minutes, 72°C for 3 minutes, and the final incubation was at 72°C for 10 minutes.

[0068] The main product of the PCR reaction was a 413 bp fragment, which was gel purified. The purified fragment was ligated to the large fragment obtained from pBR322 digested with HindIII and Sall fragment, yielding the plasmid pBR322/S2d.

[0069] Ligation of the 413 bp HindIII-Sall S2 fragment with the 1.36 kb BamHI-HindIII fragment containing the ADH2/GAP promoter, and with the large BamHI-Sall fragment of the yeast vector pBS24.1 yielded recombinant vectors, which were cloned. Correct recombinant vectors were identified by the presence of a 1.77 kb fragment after digestion with BamHI and Sall. An expression vector constructed from the amplified sequence, and containing the sequence encoding S2 fused directly to the ADH2/GAP promoter is identified as pS2d#9.

Example 4: Synthesis of HCV C Antigen

[0070] HCV antigen C22 is from the C domain. It comprises amino acids 1-122 of Figure 1.

[0071] The protocol for the construction of the expression vector encoding the C polypeptide and for its expression in yeast was analogous to that used for the expression of the C100 polypeptide, described supra, except for the following.

[0072] The template for the PCR reaction was pBR322/Ag30a which had been linearized with HindIII. Ag30 is a cDNA clone that encodes amino acids 1-122. The oligonucleotides used as primers for the amplification by PCR of the C encoding sequence were the following.
For the 5'-region of the C sequence:

5' GAG TGC AGC TTC AAA ACA AAA TGA GCA CGA
ATC CTA AAC CTC AAA AAA AAA AC 3' ,

and
for the 3'-region of the C sequence:

5' GAG TGC TCG TCG ACT CAT TAA CCC AAA TTG CGC
GAC CTA CGC CGG GGG TCT GT 3' .

The primer for the 5'-region introduces a HindIII site into the amplified product, and the primer for the 3'-region introduces translation stop codons and a Sall site. The PCR was run for 29 cycles of 94°C for a minute, 37°C for 2 minutes, 72°C for 3 minutes, and the final incubation was at 72°C for 10 minutes.

[0073] The major product of PCR amplification is a 381 bp polynucleotide. Ligation of this fragment with the Sall-HindIII large Sall-HindIII fragment of pBR322 yielded the plasmid pBR322/C2.

[0074] Ligation of the 381 bp HindIII-Sall C coding fragment excised from pBR322/C2 with the 1.36 kb BamHI-HindIII fragment containing the ADH2/GAP promoter, and with the large BamHI-Sall fragment of the yeast vector pBS24.1 yielded recombinant vectors, which were cloned. Correct recombinant vectors were identified by the presence of a 1.74 kb fragment after digestion with BamHI and Sall. An expression vector constructed from the amplified sequence, and containing the sequence encoding C fused directly to the ADH2/GAP promoter is identified as pC22.

[0075] Analysis for expressed C polypeptide by the transformed cells was performed on total cell lysates and crude extracts prepared from single yeast colonies obtained from the Leu⁻ plates. The cell lysates and crude extracts were analyzed by electrophoresis on SDS polyacrylamide gels. The C polypeptide is expected to have 122 amino acids and by gel analysis the expressed polypeptide has a MW_r of approximately 13.6 Kd.

Example 5: Synthesis of NS5 Polypeptide

[0076] This polypeptide contains sequence from the N-terminus of the NS5 domain. Specifically it includes amino acids 2054 to 2464 of Figure 1. The protocol for the construction of the expression vector encoding the NS5 polypeptide and for its expression were analogous to that used for the expression of C33c (see Example 1).

Example 6: Radioimmunoassay (RIA) for Antibodies to HCV

[0077] The HCV antigens of Examples 1-5 were tested in an RIA format for their ability to detect antibodies to HCV in the serum of individuals clinically diagnosed as having HCV (Non-A, Non-B) and in serum from blood given by paid blood donors.

[0078] The RIA was based upon the procedure of Tsu and Herzenberg (1980) in SELECTED METHODS IN CELLULAR IMMUNOLOGY (W.H. Freeman & Co.), pp. 373-391. Generally, microtiter plates (Immulon 2, Removawell strips) are coated with purified HCV antigen. The coated plates are incubated with the serum samples or appropriate controls. During incubation, antibody, if present, is immunologically bound to the solid phase antigen. After removal of the unbound material and washing of the microtiter plates, complexes of human antibody-NANBV antigen are detected by incubation with ¹²⁵I-labeled sheep anti-human immunoglobulin. Unbound labeled antibody is removed by aspiration, and the plates are washed. The radioactivity in individual wells is determined; the amount of bound human anti-HCV antibody is proportional to the radioactivity in the well.

[0079] Specifically, one hundred microliter aliquots containing 0.1 to 0.5 micrograms of the HCV antigen in 0.125 M Na borate buffer, pH 8.3, 0.075 M NaCl (BBS) was added to each well of a microliter plate (Dynatech Immulon 2 Removawell Strips). The plate was incubated at 4°C overnight in a humid chamber, after which, the antigen solution was removed and the wells washed 3 times with BBS containing 0.02% Triton X-100^(TM) (BBST). To prevent nonspecific binding, the wells were coated with bovine serum albumin (BSA) by addition of 100 microliters of a 5 mg/ml solution of BSA in BBS followed by incubation at room temperature for 1 hour; after this incubation the BSA solution was removed. The antigens in the coated wells were reacted with serum by adding 100 microliters of serum samples diluted 1:100 in 0.01M Na phosphate buffer, pH 7.2, 0.15 M NaCl (PBS) containing 10 mg/ml BSA, and incubating the serum

containing wells for 1 hr at 37°C. After incubation, the serum samples were removed by aspiration, and the wells were washed 5 times with BBST. Antibody bound to the antigen was determined by the binding of ^{125}I -labeled $\text{F}'(\text{ab})_2$ sheep anti-human IgG to the coated wells. Aliquots of 100 microliters of the labeled probe (specific activity 5-20 microcuries/microgram) were added to each well, and the plates were incubated at 37°C for 1 hour, followed by removal of excess probe by aspiration, and 5 washes with BBST. The amount of radioactivity bound in each well was determined by counting in a counter which detects gamma radiation.

[0080] Table 1 below presents the results of the tests on the serum from individuals diagnosed as having HCV.

Table 1

INDIVIDUAL	ANTIGEN				
	S2	C22	C100	C33c	NS5
CVH IVDA	P	P	P(+++)	P	P
CVH IVDA	P	P	P(+)	P	P
CVH IVDA	P	P	P(+)	P	P
CVH NOS	P	P	P	P	P
AVH NOS HS	N	N	N	N	N
AVH NOS HS	P	N	N	N	N
AVH NOS HS	P	N	N	N	N
AVH NOS HS	P/N	N	N	N	N
AVH PTVH	N	N	N	P/N	N
AVH NOS HS	N	N	N	N	N
AVH NOS	N	N	N	N	P
AVH PTVH	N	N	N	N	N
AVH IVDA	N	P	N	P	P
AVH PTVH	P	P/N	N	N	P
AVH NOS	N	P	N	N	N
AVH IVDA	N	P	N	P	P
AVH NOS HS	P/N	N	N	N	N
AVH PTVH	N	N	N	N	N
CVH IVDA	P	P	P	P	P
CVH IVDA	P	P	P	P	P
AVH NOS HS	N	N	N	N	N
CVH PTVH	P	P	N	N	N
AVH PTVH	P	N	P(+)	P(+++)	N
CVH PTVH	N	P	P	P	P
CVH NOS HS	P	P	P	P	N
CVH NOS	N	P	P/N	P	P
CVH IVDA	N	N	N	P	N
AVH IVDA	P	P	P	P	P
AVH IVDA	P	P	P	P	P
CVH IVDA	P	P	P	P	P
AVH IVDA	P/N	P	N	P	P
AVH IVDA	N	P	P	P	N
CVH PTVH	P	P/N	N	N	N
CVH NOS	N	N	N	N	N
CVH NOS	N	N	N	N	N
CVH IVDA	P	P	P	P	P
AVH IVDA	P	P	P	P	P
CVH PTVH	P	P	P	P	P
AVH PTVH?	N	P	P	P	P
AVH IVDA	N	P	N	P	N
AVH NOS	N	N	N	N	N

EP 0 693 687 B1

Table 1 (continued)

	INDIVIDUAL	ANTIGEN				
		S2	C22	C100	C33c	NS5
5	AVH NOS	N	N	N	N	N
	CVH NOS	N	P	N	N	P
	CVH NOS	P	P	N	N	N
	CVH NOS HS	P	P	P	P	P
10	CVH PTVH	P	P	N	P	P
	AVH nurse	P	P	N	N	N
	AVH IVDA	P	P	P	P	N
	AVH IVDA	N	P	P(+)	P(+++)	N
	AVH nurse	P/N	P	N	N	N
15	AVH PTVH	P/N	P	P	N	P
	AVH NOS	N	P/N	N	N	P
	AVH NOS	N	P	N	P	N
	AVH PTVH	P	P/N	N	N	N
20	AVH PTVH	N	P	N	P	P
	AVH PTVH	P	P	P	P	P
	AVH PTVH	N	P	N	N	P
	CVH PTVH	P/N	P	P(+)	P(+++)	N
	AVH PTVH	N	P/N	P(+)	P(+++)	P
25	AVH PTVH	P	(?)	P	N	P
	CVH PTVH	N	P	N	P	P
	CVH PTVH	N	P	P	P	P
	CVH PTVH	N	N	N	N	N
30	AVH NOS	N	N	N	N	N
	AVH nurse	P	P	N	N	N
	CVH PTVH	N	P	N	N	P
	AVH IVDA	N	P	N	P/N	N
	CVH PTVH	P	P	P(+)	P(+++)	P
35	AVH NOS	P	P	N	N	N
	AVH NOS	P/N	P	N	N	P
	AVH PTVH	P/N	P	P	P	P
	AVH NOS	N	P	P	P	P/N
40	AVH IVDA	N	P	N	N	P
	AVH NOS	N	P/N	N	N	N
	AVH NOS	P	P	N	N	P
	AVH PTVH	N	P	P	P	P
	crypto	P	P	P	P	P
45	CVH NOS	N	P	P	P	P
	CVH NOS	N	N	N	N	N
	AVH PTVH	N	P	P(+)	P(++)	N
	AVH PTVH	N	P/N	P(+)	P(++)	P
50	AVH PTVH	N	P/N	P(+)	P(++)	P
	CVH IVDA	P	P	P	P	P
	CVH IVDA	P	P	P	P	P
	CVH IVDA	P	P	P	P	P
	CVH IVDA	P	P	P	P	P
55	AVH NOS	N	P	N	N	N
	CVH IVDA	P	P	P	P	P/N
	AVH IVDA	P	P	P	P	N

EP 0 693 687 B1

Table 1 (continued)

	INDIVIDUAL	ANTIGEN				
		S2	C22	C100	C33c	NS5
5	AVH NOS	P	P	N	N	N
	AVH NOS	P	P	N	N	N
	CVH PTVH	P	P	N	N	P/N
	AVH PTVH	N	P	N	P	P
10	AVH NOS	N	N	N	N	N
	AVH NOS	N	P	N	N	N
	AVH NOS	P	N	N	N	N
	CVH NOS	N	N	N	N	N
	AVH NOS	N	P/N	N	N	N
15	AVH IVDA	N	P	P	P	P
	crypto	N	P	N	N	P/N
	crypto	P	P	P/N	P	P
	AVH IVDA	N	N	P	P	N
20	AVH IVDA	N	P	P	P	N
	AVH NOS	N	N	N	N	N
	AVH NOS	N	N	N	N	N
	CVH IVDA	P	P	P	P	P
	CVH PTVH	N	N	N	N	N
25	CVH PTVH	P	P	P(+)	P(+++)	P
	CVH PTVH	P	P	P(+)	P(+++)	P
	CVH NOS	P/N	N	N	N	N
	CVH NOS	N	N	N	N	N
30	CVH PTVH	P	P	P	P	P
	CVH PTVH	P	P	P	P	P
	CVH PTVH	P	P	P	P	P
	AVH IVDA	N	P	P	P	P
	CVH NOS	N	N	N	N	N
35	CVH NOS	N	N	N	N	N
	CVH PTVH	P	P	P	P	P
	AVH NOS	P	P	N	N	P/N
	AVH NOS	N	P/N	N	N	N
40	CVH PTVH	P	P	N	N	P
	CVH NOS	N	P/N	N	N	N
	AVH NOS	N	P	N	N	N
	AVH NOS	N	P	N	N	N
	CVH PTVH	N	P	N	N	N
45	AVH IVDA	N	P	N	P	P
	AVH NOS	P	N	N	N	N
	CVH NOS	N	N	N	N	N
	CVH NOS	N	N	N	N	N
50	CVH IVDA	P	P	P	P	P
	CVH IVDA	P/N	P	P	P	P
	CVH IVDA	P	P	P	P	P
	CVH IVDA	N	P	P	P	P
	AVH NOS	N	P	N	N	N
55	CVH IVDA	N	P	N	N	P
	CVH IVDA	N	P	N	N	P
	AVH PTVH	P	P	N	P	P

EP 0 693 687 B1

Table 1 (continued)

	INDIVIDUAL	ANTIGEN				
		S2	C22	C100	C33c	NS5
5	AVH PTVH	P	P	N	P	P
	CVH NOS	N	P/N	N	N	P/N
	CVH NOS	N	P	N	N	N
	CVH NOS	N	N	N	N	N
10	CVH PTVH	P	P	P	P	P
	CVH PTVH	P	P	P	P	P
	CVH PTVH	P	P	P	P	P
	AVH IVDA	N	P	N	N	P
	AVH IVDA	N	P	P(++)	P(+)	P
15	CVH PTVH	P	P	P	P	P
	AVH PTVH	N	P	P	P	P
	CVH PTVH?	N	P	P	P	P
	CVH PTVH?	P/N	P	P	P	P
20	CVH NOS HS	P	P	N	N	N
	CVH IVDA	P	P	P	P	N
	CVH PTVH	N	P	P	P	P
	CVH PTVH	P	P	P	P	P/N
	CVH NOS	P	P	P	P	P
25	CVH IVDA	P	P	P	P	P
	CVH PTVH	P	P	P	P	N
	CVH PTVH	P	P	P	P	P
	CVH NOS	N	N	N	N	P/N
30	CVH NOS	N	P/N	N	N	P/N
	CVH PTVH	P	P	P	P	P
	CVH NOS	N	P	N	N	N
	CVH NOS	N	N	N	N	N
	CVH NOS	P	P	N	N	P/N
35	CVH NOS	N	N	N	N	N
	CVH NOS HS	P	P	P	P	P
	CVH NOS HS	P	P	P	P	P
	CVH PTVH	N	N	N	N	N
40	AVH PTVH	N	P	P	P	P
	AVH NOS			-	-	
	CVH PTVH	N	P	P/N	P(+++)	N
	crypto	P	P	P	P	P
	crypto	P	P	P	P	P
45	crypto	N	P	N	N	N
	crypto	N	P	P	P	P
	CVH PTVH	P	P	P	P	P
	crypto	N	N	N	N	N
	crypto	N	P	N	N	P/N
50	crypto	N	P	N	N	P
	crypto	P	P	P	P	P
	crypto	N	P	N	P	N
	crypto			-	-	
55	crypto			-	-	
	CVH NOS			-	-	
	AVH-IVDA	N	P	N	P(+)	P

EP 0 693 687 B1

Table 1 (continued)

INDIVIDUAL	ANTIGEN				
	S2	C22	C100	C33c	NS5
AVH-IVDA	N	P/N	N	P(++)	N
AVH = acute viral hepatitis CVH = chronic viral hepatitis PTVH = post-transfusion viral hepatitis IVDA = intravenous drug abuser crypto = cryptogenic hepatitis NOS = non-obvious source P = positive N = negative					

[0081] Per these results, no single antigen reacted with all sera. C22 and C33c were the most reactive and S2 reacted with some sera from some putative acute HCV cases with which no other antigen reacted. Based on these results, the combination of two antigens that would provide the greatest range of detection is C22 and C33c. If one wished to detect a maximum of acute infections, S2 would be included in the combination.

[0082] Table 2 below presents the results of the testing on the paid blood donors.

Table 2

Donor	Antigens				
	C100	C33c	C22	S2	NS5
1	N	N	N	N	N
2	N	N	N	N	N
3	P	P	P	P	P
4	N	N	N	N	N
5	N	N	N	N	N
6	N	N	N	N	N
7	N	N	N	N	N
8	N	N	N	N	N
9	N	N	N	N	N
10	N	N	N	N	N
11	N	N	N	N	N
12	N	N	N	N	N
13	N	N	N	N	N
14	N	N	N	N	N
15	N	N	N	N	N
16	N	N	N	N	N
17	N	N	N	N	N
18	P	P	P	P	P
19	P	P	N	P	P
20	P	P	N	P	P
21	N	N	N	N	N
22	N	P	P	N	P
23	P	P	P	P	P
24	N	N	N	N	N
25	N	N	N	N	N
26	N	N	N	N	N
27	N	N	N	N	N
28	N	N	N	N	N
29	N	N	N	N	N

Table 2 (continued)

Donor	Antigens				
	C100	C33c	C22	S2	NS5
30	N	N	N	N	N
31	P	P	P	N	P
32	N	N	N	N	N
33	N	N	N	N	N
34	N	N	N	N	P
35	N	N	P	N	P
36	N	N	N	N	N
37	N	N	N	N	N
38	N	N	N	N	N
39	N	N	N	N	N
40	N	N	N	N	N
41	N	N	N	N	P
42	N	N	N	N	N
43	N	N	N	N	N
44	N	N	N	N	N
45	N	N	N	N	N
46	N	N	N	N	N
47	P	P	N	N	P
48	N	N	N	N	N
49	N	N	N	N	N
50	N	N	N	N	N
51	N	P	P	N	P
52	N	N	N	N	N
53	N	P	P	N	P
54	P	P	P	P	N
55	N	N	N	N	N
56	N	N	N	N	N
57	N	N	N	N	N
58	N	N	N	N	N
59	N	N	N	N	N
60	N	N	N	N	N
61	N	N	N	N	N
62	N	N	N	N	N
63	N	N	N	N	N
64	N	N	N	N	N
65	N	N	N	N	N
66	N	N	N	N	N
67	N	N	N	N	N
68	N	N	N	N	N
69	N	N	N	N	N
70	P	P	P	P	P
71	N	N	N	N	N
72	N	N	N	N	N
73	P	P	P	P	N
74	N	N	N	N	N
75	N	N	N	N	N
76	N	N	N	N	P
77	N	N	N	N	N

EP 0 693 687 B1

Table 2 (continued)

	Donor	Antigens				
		C100	C33c	C22	S2	NS5
5	78	N	N	N	N	N
	79	N	N	N	N	N
	80	N	N	N	N	N
	81	N	N	N	N	N
10	82	N	N	N	N	N
	83	P	P	N	N	N
	84	N	N	P	N	N
	85	N	N	N	N	N
15	86	P	P	P	P	N
	87	N	N	N	N	N
	88	N	N	N	N	N
	89	P	P	P	P	P
	90	P	P	P	P	N
20	91	N	N	N	N	P
	92	P	P	P	N	N
	93	N	N	N	N	N
	94	N	N	N	N	N
	95	N	N	N	N	N
25	96	N	N	N	N	N
	97	N	N	N	N	N
	98	N	P	P	P	P
	99	P	P	P	P	P
30	100	N	N	N	N	N
	101	P	P	P	P	P
	102	N	N	N	N	N
	103	N	N	N	N	N
	104	N	N	N	N	N
35	105	P	P	P	P	N
	106	N	N	N	N	N
	107	N	N	N	N	N
	108	N	N	N	N	N
40	109	P	P	P	P	P
	110	P	P	P	N	P
	111	P	P	P	N	P
	112	N	N	N	N	N
45	113	P	P	P	P	P
	114	N	N	N	N	N
	115	N	N	N	N	N
	116	P	P	P	P	P
	117	N	N	N	N	N
50	118	N	N	N	N	N
	119	N	N	N	N	N
	120	P	P	P	P	P
	121	N	N	N	N	N
55	122	N	P	P	N	P
	123	N	N	N	N	N
	124	N	N	N	N	N
	125	N	N	N	N	N

EP 0 693 687 B1

Table 2 (continued)

Donor	Antigens				
	C100	C33c	C22	S2	NS5
126	P	N	N	N	N
127	N	N	N	N	N
128	N	N	N	N	N
129	N	N	N	N	N
130	P	P	P	P	N
131	N	N	N	N	P
132	N	N	N	N	N
133	N	N	N	N	N
134	N	N	N	N	N
135	N	N	N	N	N
136	N	N	N	N	N
137	N	N	N	N	N
138	N	N	N	N	N
139	N	N	N	N	N
140	P	N	N	N	N
141	P	N	P	P	P
142	N	N	N	N	N
143	N	N	N	N	N
144	N	N	N	N	N
145	N	N	N	N	N
146	N	N	N	N	N
147	N	N	N	N	N
148	N	N	N	N	N
149	N	N	N	N	N
150	N	N	N	N	N
151	N	N	N	N	N
152	N	N	N	N	N
153	N	N	N	N	N
154	P	P	P	P	P
155	N	N	N	N	N
156	N	N	N	N	N
157	N	N	N	N	N
158	N	N	N	N	N
159	N	N	N	N	N
160	N	N	N	N	N
161	P	P	P	P	P
162	N	N	N	N	N
163	N	N	N	N	N
164	P	P	P	N	P
165	N	N	N	N	N
166	P	P	P	N	P
167	N	N	N	N	N
168	N	N	N	N	N
169	N	N	N	N	N
170	N	N	N	N	N
171	N	N	N	N	N
172	N	N	N	N	N
173	N	N	N	N	N

Table 2 (continued)

Donor	Antigens				
	C100	C33c	C22	S2	NS5
174	N	N	N	N	N
175	N	N	N	N	N
176	N	N	N	N	N
177	N	N	N	N	P
178	N	N	N	N	N
179	N	N	N	N	N
180	N	N	N	N	N
181	N	N	N	N	N
182	N	N	N	N	N
183	P	P	P	P	P
184	N	N	N	N	N
185	N	N	N	N	N
186	N	N	N	N	N
187	N	N	N	N	N
188	N	P	P	N	N
189	N	N	N	N	N
190	N	N	N	N	N
191	N	N	N	N	N
192	N	N	N	N	N
193	N	N	N	N	N
194	N	N	N	N	N
195	N	N	N	N	N
196	N	N	N	N	N
197	N	N	N	N	P
198	P	P	P	N	N
199	N	N	N	N	P
200	P	P	P	P	N

[0083] The results on the paid donors generally confirms the results from the sera of infected individuals.

Example 7: ELISA Determinations of HCV Antibodies Using Combination of HCV Antigens

[0084] Plates coated with a combination of C22 and C33c antigens are prepared as follows. A solution containing coating buffer (50mM Na Borate, pH 9.0), 21 ml/plate, BSA (25 micrograms/ml), C22 and C33c (2.50 micrograms/ml each) is prepared just prior to addition to the Removeawell Immulon I plates (Dynatech Corp.). After mixing for 5 minutes, 0.2ml/well of the solution is added to the plates, they are covered and incubated for 2 hours at 37°C, after which the solution is removed by aspiration. The wells are washed once with 400 microliters wash buffer (100 mM sodium phosphate, pH 7.4, 140 mM sodium chloride, 0.1% (W/V) casein, 1% (W/V) Triton x-100TM, 0.01% (W/V) Thimerosal). After removal of the wash solution, 200 microliters/well of Postcoat solution (10 mM sodium phosphate, pH 7.2, 150 mM sodium chloride, 0.1% (w/v) casein, 3% sucrose and 2 mM phenylmethylsulfonylfluoride (PMSF)) is added, the plates are loosely covered to prevent evaporation, and are allowed to stand at room temperature for 30 minutes. The wells are then aspirated to remove the solution, and lyophilized dry overnight, without shelf heating. The prepared plates may be stored at 2-8°C in sealed aluminum pouches with dessicant (3 g Sorb-itTM packs).

[0085] In order to perform the ELISA determination, 20 microliters of serum sample or control sample is added to a well containing 200 microliters of sample diluent (100 mM sodium phosphate, pH 7.4, 500 mM sodium chloride, 1 mM EDTA, 0.1% (W/V) Casein, 0.01% (W/V) Thimerosal, 1% (W/V) Triton X-100TM, 100 micrograms/ml yeast extract). The plates are sealed, and are incubated at 37°C for two hours, after which the solution is removed by aspiration, and the wells are washed three times with 400 microliters of wash buffer (phosphate buffered saline (PBS) containing 0.05% Tween 20TM). The washed wells are treated with 200 microliters of mouse anti-human IgG-horse radish peroxidase (HRP) conjugate contained in a solution of Ortho conjugate diluent (10 mM sodium phosphate, pH 7.2, 150 mM sodium

chloride, 50% (v/v) fetal bovine serum, 1% (V/V) heat treated horse serum, 1 mM $K_3Fe(CN)_6$, 0.05% (W/V) Tween 20, 0.02% (W/V) Thimerosal). Treatment is for 1 hour at 37°C, the solution is removed by aspiration, and the wells are washed three times with 400 ml wash buffer, which is also removed by aspiration. To determine the amount of bound enzyme conjugate, 200 microliters of substrate solution (10 mg O-phenylenediamine dihydrochloride per 5 ml of Developer solution) is added. Developer solution contains 50 mM sodium citrate adjusted to pH 5.1 with phosphoric acid, and 0.6 microliters/ml of 30% H_2O_2 . The plates containing the substrate solution are incubated in the dark for 30 minutes at room temperature, the reactions are stopped by the addition of 50 microliters/ml 4N sulfuric acid, and the ODs determined.

[0086] In a similar manner, ELISAs using fusion proteins of C22 and C33c, and C22, C33c, and S2 and combinations of C22 and C100, C22 and S2, C22 and an NS5 antigen, C22, C33c, and S2, and C22, C100, and S2 may be carried out.

[0087] Modifications of the above-described modes for carrying out the invention that are obvious to those of skill in the fields of molecular biology, immunology, and related fields are intended to be within the scope of the following claims.

Claims

Claims for the following Contracting States : AT, BE, CH, LI, DE, DK, FR, GB, GR, IT, LU, NL, SE

1. A combination of hepatitis C virus (HCV) antigens in one or more polypeptides made by chemical synthesis or recombinant expression, comprising:

- (a) a first HCV antigen comprising an epitope from the C domain of the HCV polyprotein; and
- (b) a second HCV antigen selected from

- (i) an HCV antigen comprising an epitope from the S domain;
- (ii) an HCV antigen comprising an epitope from the NS3 domain;
- (iii) an HCV antigen comprising an epitope from the NS4 domain; and
- (iv) an HCV antigen comprising an epitope from the NS5 domain;

with the proviso that the combination is not the peptide p1 with C100-3, the peptide p35 with C100-3, the peptide p99 with C100-3 or a peptide having amino acids 9 to 177 of the HCV-1 polyprotein.

2. A combination according to claim 1 wherein said second HCV antigen comprises an epitope from the S domain and which combination further comprises at least one additional HCV antigen selected from:

- (i) an HCV antigen comprising an epitope from the NS3 domain;
- (ii) an HCV antigen comprising an epitope from the NS4 domain; and
- (iii) an HCV antigen comprising an epitope from the NS5 domain;

3. A combination according to claim 2 wherein the additional antigen is from the NS3 domain.

4. A combination according to claim 3 wherein the first HCV antigen is C22 and the additional HCV antigen is C33c.

5. A combination according to claim 2 wherein the additional antigen is from the NS4 domain.

6. A combination according to claim 5 wherein the first HCV antigen is C22 and the additional HCV antigen is C100.

7. A combination according to any one of claims 1 to 6 wherein the antigen from the S domain is antigen S2.

8. A combination according to any one of claims 1 to 7 wherein the combination is in the form of a fusion polypeptide.

9. A combination according to any one of claims 1 to 8 wherein the combination is in the form of said first HCV antigen and said additional antigens bound to a common solid matrix.

10. A combination according to claim 9 wherein the solid matrix is the surface of a microtiter plate well, a bead or a dipstick.

11. A combination according to any one of claims 1 to 7 wherein the combination is immobilised on the surface of a solid matrix suitable for detection of HCV by immunoassay.
12. A method for detecting antibodies to HCV in a mammalian body component suspected of containing said antibodies comprising contacting said body component with a combination of antigens according to any one of claims 1 to 11 under conditions that permit antibody antigen reaction and detecting the presence of immune complexes of said antibodies and said antigens.
13. A kit for carrying out an assay for detecting antibodies to hepatitis C antigen (HCV) in a mammalian body component suspected of containing said antibodies comprising in packaged combination:
 - (a) the combination according to any one of claims 1 to 11;
 - (b) standard control reagents; and
 - (c) instructions for carrying out the assay.

Claims for the following Contracting State : ES

1. A method for detecting antibodies to hepatitis C virus (HCV) in a mammalian body component suspected of containing said antibodies comprising contacting said body component with a combination of HCV antigens under conditions that permit antibody-antigen reaction and detecting the presence of immune complexes of said antibodies and said antigens, wherein the antigens are in one or more polypeptides made by chemical synthesis or recombinant expression, immobilized on the surface of a solid matrix suitable for detection of HCV by immunoassay, comprising:
 - (a) a first HCV antigen comprising an epitope from the C domain of the HCV polyprotein; and
 - (b) a second HCV antigen selected from
 - (i) an HCV antigen comprising an epitope from the S domain;
 - (ii) an HCV antigen comprising an epitope from the NS3 domain;
 - (iii) an HCV antigen comprising an epitope from the NS4 domain; and
 - (iv) an HCV antigen comprising an epitope from the NS5 domain;with the proviso that the combination is not the peptide p1 with C100-3, the peptide p35 with C100-3 the peptide p99 with C100-3 or a peptide having amino acids 9 to 177 of the HCV-1 polyprotein.
2. A method according to claim 1 wherein said second HCV antigen comprises an epitope from the S domain and which combination further comprises at least one additional HCV antigen selected from:
 - (i) an HCV antigen comprising an epitope from the NS3 domain;
 - (ii) an HCV antigen comprising an epitope from the NS4 domain; and
 - (iii) an HCV antigen comprising an epitope from the NS5 domain;
3. A method according to claim 2 wherein the additional antigen is from the NS3 domain.
4. A method according to claim 3 wherein the first HCV antigen is C22 and the additional HCV antigen is C33c.
5. A method according to claim 2 wherein the additional antigen is from the NS4 domain.
6. A method according to claim 5 wherein the first HCV antigen is C22 and the additional HCV antigen is C100.
7. A method according to any one of claims 1 to 6 wherein the antigen from the S domain is antigen S2.
8. A method according to any one of claims 1 to 7 wherein the combination is in the form of a fusion polypeptide.
9. A method according to any one of claims 1 to 8 wherein the combination is in the form of said first HCV antigen and said additional antigens bound to a common solid matrix.

10. A method according to claim 9 wherein the solid matrix is the surface of a microtiter plate well, a bead or a dipstick.
11. A combination of hepatitis C virus (HCV) antigens in one or more polypeptides made by chemical synthesis or recombinant expression, comprising:
 - (a) a first HCV antigen comprising an epitope from the C domain of the HCV polyprotein; and
 - (b) a second HCV antigen selected from
 - (i) an HCV antigen comprising an epitope from the S domain;
 - (ii) an HCV antigen comprising an epitope from the NS3 domain;
 - (iii) an HCV antigen comprising an epitope from the NS4 domain; and
 - (iv) an HCV antigen comprising an epitope from the NS5 domain;with the proviso that the combination is not the peptide p1 with C100-3, the peptide p35 with C100-3, the peptide p99 with C100-3 or a peptide having amino acids 9 to 177 of the HCV-1 polyprotein.
12. A combination according to claim 12 wherein said second HCV antigen comprises an epitope from the S domain and which combination further comprises at least one additional HCV antigen selected from:
 - (i) an HCV antigen comprising an epitope from the NS3 domain;
 - (ii) an HCV antigen comprising an epitope from the NS4 domain; and
 - (iii) an HCV antigen comprising an epitope from the NS5 domain;
13. A combination according to claim 12 wherein the additional antigen is from the NS3 domain.
14. A combination according to claim 13 wherein the first HCV antigen is C22 and the additional HCV antigen is C33c.
15. A combination according to claim 14 wherein the additional antigen is from the NS4 domain.
16. A combination according to claim 15 wherein the first HCV antigen is C22 and the additional HCV antigen is C100.
17. A combination according to any one of claims 11 to 16 wherein the antigen from the S domain is antigen S2.
18. A combination according to any one of claims 11 to 17 wherein the combination is in the form of a fusion polypeptide.
19. A combination according to any one of claims 11 to 18 wherein the combination is in the form of said first HCV antigen and said additional antigens bound to a common solid matrix.
20. A combination according to claim 19 wherein the solid matrix is the surface of a microtiter plate well, a bead or a dipstick.
21. A combination according to any one of claims 11 to 20 wherein the combination is immobilised on the surface of a solid matrix suitable for detection of HCV by immunoassay.
22. A kit for carrying out an assay for detecting antibodies to hepatitis C antigen (HCV) in a mammalian body component suspected of containing said antibodies comprising in packaged combination:
 - (a) the combination according to any one of claims 11 to 21;
 - (b) standard control reagents; and
 - (c) instructions for carrying out the assay.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, LI, DE, DK, FR, GB, GR, IT, LU, NL, SE

1. Kombination von Hepatitis-C-Virus(HCV)-Antigenen in einem Polypeptid oder in mehreren Polypeptiden, das/die

durch chemische Synthese oder rekombinante Expression hergestellt wurde/wurden, umfassend :

- (a) ein erstes HCV-Antigen, das ein Epitop von der C-Domäne des HCV-Polyproteins enthält; und
 - (b) ein zweites HCV-Antigen, ausgewählt aus
 - (i) einem HCV-Antigen, das ein Epitop von der S-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (iii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iv) ein HCV-Antigen, das ein Epitop von der NS5-Domäne enthält;
- mit der Maßgabe, daß die Kombination nicht das Peptid p1 mit C100-3, das Peptid p35 mit C100-3, das Peptid p99 mit C100-3 oder einem Peptid mit den Aminosäuren 9 bis 177 des HCV-1-Polyproteins ist.
2. Kombination nach Anspruch 1, wobei das zweite HCV-Antigen ein Epitop von der S-Domäne enthält und wobei die Kombination weiterhin wenigstens ein weiteres HCV-Antigen, ausgewählt aus:
 - (i) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iii) einem HCV-Antigen, das ein Epitop von der NS5-Domäne enthält;enthält.
3. Kombination nach Anspruch 2, wobei das zusätzliche Antigen von der NS3-Domäne ist.
4. Kombination nach Anspruch 3, wobei das erste HCV-Antigen C22 ist und das zusätzliche HCV-Antigen C33c ist.
5. Kombination nach Anspruch 2, wobei das zusätzliche Antigen von der NS4-Domäne ist.
6. Kombination nach Anspruch 5, wobei das erste HCV-Antigen C22 ist und das zusätzliche HCV-Antigen C100 ist.
7. Kombination nach einem der Ansprüche 1 bis 6, wobei das Antigen von der S-Domäne das Antigen S2 ist.
8. Kombination nach einem der Ansprüche 1 bis 7, wobei die Kombination in Form eines Fusionspolypeptids ist.
9. Kombination nach einem der Ansprüche 1 bis 8, wobei die Kombination in Form des genannten ersten HCV-Antigens und der genannten zusätzlichen Antigene, die an eine gemeinsame feste Matrix gebunden sind, ist.
10. Kombination nach Anspruch 9, wobei die feste Matrix die Oberfläche einer Mikrotiterplattenvertiefung, einer Kugel oder eines Eintauchstäbchens ist.
11. Kombination nach einem der Ansprüche 1 bis 7, wobei die Kombination auf der Oberfläche einer festen Matrix, die zum Nachweis von HCV durch Immunoassay geeignet ist, immobilisiert ist.
12. Verfahren zum Nachweis von Antikörpern gegen HCV in einem Säugetierkörperbestandteil, der im Verdacht steht, diese Antikörper zu enthalten, umfassend das Inkontaktbringen des Körperbestandteils mit einer Kombination von Antigenen nach einem der Ansprüche 1 bis 11 unter Bedingungen, die die Antikörper-Antigen-Reaktion ermöglichen, und Nachweis des Vorhandenseins von Immunkomplexen der genannten Antikörper und der genannten Antigene.
13. Kit zum Durchführen eines Assays zum Nachweis von Antikörpern gegen Hepatitis-C-Antigen (HCV) in einem Säugetierkörperbestandteil, der im Verdacht steht, diese Antikörper zu enthalten, umfassend:
 - (a) die Kombination nach einem der Ansprüche 1 bis 11;
 - (b) Standardkontrollreagentien; und
 - (c) Anleitungen zum Durchführen des Assays;in verpackter Kombination.

Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zum Nachweis von Antikörpern gegen das Hepatitis-C-Virus (HCV) in einem Säugetierkörperbestandteil, der im Verdacht steht, die genannten Antikörper zu enthalten, umfassend das Inkontaktbringen des Körperbestandteils mit einer Kombination von HCV-Antigenen unter Bedingungen, die eine Antikörper-Antigen-Reaktion ermöglichen, und Nachweis des Vorhandenseins von Immunkomplexen der genannten Antikörper und der genannten Antigene, wobei die Antigene in einem Polypeptid oder in mehreren Polypeptiden sind, das/die durch chemische Synthese oder rekombinante Expression hergestellt wurde/wurden, auf der Oberfläche einer festen Matrix, die zum Nachweis von HCV durch Immunoassay geeignet sind, immobilisiert wurde/wurden, umfassend:
 - (a) ein erstes HCV-Antigen, das ein Epitop von der C-Domäne des HCV-Polypeptids enthält; und
 - (b) ein zweites HCV-Antigen, ausgewählt aus
 - (i) einem HCV-Antigen, das ein Epitop von der S-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (iii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iv) einem HCV-Antigen, das ein Epitop von der NS5-Domäne enthält;mit der Maßgabe, daß die Kombination nicht das Peptid p1 mit C100-3, das Peptid p35 mit der C100-3, das Peptid p99 mit C100-3 oder einem Peptid, das die Aminosäuren 9 bis 177 des HCV-1-Polypeptids besitzt, ist.
2. Verfahren nach Anspruch 1, wobei das zweite HCV-Antigen ein Epitop von der S-Domäne enthält und wobei die Kombination weiterhin wenigstens ein zusätzliches HCV-Antigen, ausgewählt aus:
 - (i) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iii) einem HCV-Antigen, das ein Epitop von der NS5-Domäne enthält;umfaßt.
3. Verfahren nach Anspruch 2, wobei das zusätzliche Antigen von der NS3-Domäne ist.
4. Verfahren nach Anspruch 3, wobei das erste HCV-Antigen C22 ist und das zusätzliche HCV-Antigen C33c ist.
5. Verfahren nach Anspruch 2, wobei das zusätzliche Antigen von der NS4-Domäne ist.
6. Verfahren nach Anspruch 5, wobei das erste HCV-Antigen C22 ist und das zusätzliche HCV-Antigen C100 ist.
7. Verfahren nach einem der Ansprüche 1 bis 6, wobei das Antigen aus der S-Domäne das S2-Antigen ist.
8. Verfahren nach einem der Ansprüche 1 bis 7, wobei die Kombination in Form eines Fusionspolypeptids ist.
9. Verfahren nach einem der Ansprüche 1 bis 8, wobei die Kombination in der Form des ersten HCV-Antigens und der genannten zusätzlichen Antigene, die an eine gemeinsame feste Matrix gebunden sind, ist.
10. Verfahren nach Anspruch 9, wobei die feste Matrix die Oberfläche einer Mikrotiterplattenvertiefung, einer Kugel oder eines Eintauchstäbchens ist.
11. Kombination von Hepatitis-C-Virus(HCV)-Antigenen in einem oder mehreren Polypeptiden, das/die durch chemische Synthese oder rekombinante Expression hergestellt wurde/wurden, umfassend:
 - (a) ein erstes HCV-Antigen, das ein Epitop von der C-Domäne des HCV-Polypeptids enthält; und
 - (b) ein zweites HCV-Antigen, ausgewählt aus
 - (i) einem HCV-Antigen, das ein Epitop von der S-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (iii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iv) einem HCV-Antigen, das ein Epitop von der NS5-Domäne enthält;

mit der Maßgabe, daß die Kombination nicht das Peptid p1 mit C100-3, das Peptid p35 mit C100-3, das Peptid p99 mit C100-3 oder einem Peptid, das die Aminosäuren 9 bis 177 des HCV-1-Polyproteins besitzt, ist.

- 5 12. Kombination nach Anspruch 12, wobei das genannte zweite HCV-Antigen ein Epitop von der S-Domäne enthält und wobei die Kombination weiterhin wenigstens ein zusätzliches HCV-Antigen besitzt, ausgewählt aus
 - (i) einem HCV-Antigen, das ein Epitop von der NS3-Domäne enthält;
 - (ii) einem HCV-Antigen, das ein Epitop von der NS4-Domäne enthält; und
 - (iii) einem HCV-Antigen, das ein Epitop von der NS5-Domäne enthält.
- 10 13. Kombination nach Anspruch 12, wobei das zusätzliche Antigen von der NS3-Domäne ist.
14. Kombination nach Anspruch 13, wobei das erste HCV-Antigen C22 ist und das zusätzliche HCV-Antigen C33c ist.
- 15 15. Kombination nach Anspruch 14, wobei das zusätzliche Antigen von der NS4-Domäne ist.
16. Kombination nach Anspruch 15, wobei das erste HCV-Antigen C22 und das zusätzliche HCV-Antigen C100 ist.
- 20 17. Kombination nach einem der Ansprüche 11 bis 16, wobei das Antigen von der S-Domäne das S2-Antigen ist.
18. Kombination nach einem der Ansprüche 11 bis 17, wobei die Kombination in Form eines Fusionspolypeptids ist.
19. Kombination nach einem der Ansprüche 11 bis 18, wobei die Kombination in Form des genannten ersten HCV-Antigens und der genannten zusätzlichen Antigene, die an eine gemeinsame feste Matrix gebunden sind, ist.
- 25 20. Kombination nach Anspruch 19, wobei die feste Matrix die Oberfläche einer Mikrotiterplattenvertiefung, einer Kugel oder eines Eintauchstäbchens ist.
- 30 21. Kombination nach einem der Ansprüche 11 bis 20, wobei die Kombination auf der Oberfläche einer festen Matrix, die zum Nachweis von HCV durch Tmmunoassay geeignet ist, immobilisiert ist.
22. Kit zum Durchführen eines Assays zum Nachweis von Antikörpern gegen Hepatitis-C-Antigen (HCV) in einem Säugetierkörperbestandteil, der im Verdacht steht, die genannten Antikörper zu enthalten, umfassend:
 - 35 (a) die Kombination nach einem der Ansprüche 11 bis 21;
 - (b) Standardkontrollreagentien; und
 - (c) Anleitungen zum Durchführen des Assays;
- 40 in verpackter Kombination.

Revendications

45 **Revendications pour les Etats contractants suivants : AT, BE, CH, LI, DE, DK, FR, GB, GR, IT, LU, NL, SE**

1. Combinaison d'antigènes du virus de l'hépatite C (HCV) en un ou plusieurs polypeptides obtenus par synthèse chimique ou par expression recombinante, comprenant :
 - 50 (a) un premier antigène de HCV comprenant un épitope du domaine C de la polyprotéine de HCV ; et
 - (b) un second antigène de HCV choisi parmi
 - (i) un antigène de HCV comprenant un épitope du domaine S ;
 - (ii) un antigène de HCV comprenant un épitope du domaine NS3 ;
 - 55 (iii) un antigène de HCV comprenant un épitope du domaine NS4 ; et
 - (iv) un antigène de HCV comprenant un épitope du domaine NS5 ;

à la condition que la combinaison ne soit pas le peptide p1 avec C100-3, le peptide p35 avec C100-3, le

peptide p99 avec C100-3 ou un peptide ayant les acides aminés 9 à 177 de la polyprotéine de HCV-1.

2. Combinaison selon la revendication 1 dans laquelle ledit second antigène de HCV comprend un épitope du domaine S et laquelle combinaison comprend en outre au moins un antigène de HCV supplémentaire choisi parmi :
5 (i) un antigène de HCV comprenant un épitope du domaine NS3 ;
 (ii) un antigène de HCV comprenant un épitope du domaine NS4 ;
 (iii) un antigène de HCV comprenant un épitope du domaine NS5 ;
- 10 3. Combinaison selon la revendication 2 dans laquelle l'antigène supplémentaire provient du domaine NS3.
4. Combinaison selon la revendications 3 dans laquelle le premier antigène de HCV est C22 et l'antigène supplémentaire de HCV est C33c.
- 15 5. Combinaison selon la revendication 2 dans laquelle l'antigène supplémentaire provient du domaine NS4.
6. Combinaison selon la revendication 5 dans laquelle le premier antigène de HCV est C22 et l'antigène supplémentaire est C100.
- 20 7. Combinaison selon l'une quelconque des revendications 1 à 6 dans laquelle l'antigène du domaine S est l'antigène S2.
8. Combinaison selon l'une quelconque des revendications 1 à 7 dans laquelle la combinaison est sous la forme d'un polypeptide de fusion.
- 25 9. Combinaison selon l'une quelconque des revendications 1 à 8 dans laquelle la combinaison est sous la forme dudit premier antigène de HCV et lesdits antigènes supplémentaires sont liés à une matrice solide commune.
- 30 10. Combinaison selon la revendication 9 dans laquelle la matrice solide est la surface d'un puits d'une plaque de microtitration, une bille ou une bandelette.
11. Combinaison selon l'une quelconque des revendications 1 à 7, dans laquelle la combinaison est immobilisée à la surface d'une matrice solide convenant pour la détection de HCV par un test immunologique.
- 35 12. Méthode de détection des anticorps dirigés contre HCV dans un constituant corporel de mammifère suspecté de contenir lesdits anticorps comprenant la mise en contact dudit constituant corporel avec une combinaison d'antigènes selon l'une quelconque des revendications 1 à 11 dans des conditions qui permettent les réactions anticorps - antigènes et la détection de la présence de complexes immuns desdits anticorps et desdits antigènes.
- 40 13. Kit pour réaliser un test de détection d'anticorps dirigés contre un antigène de l'hépatite C (HCV) dans un constituant corporel de mammifère suspecté de contenir lesdits anticorps comprenant, sous forme de combinaison conditionnée :
45 (i) la combinaison selon l'une quelconque des revendications 1 à 11 ;
 (ii) les réactifs témoins standard ; et
 (iii) les instructions pour réaliser le test.

Revendications pour l'Etat contractant suivant : ES

- 50 1. Méthode permettant de détecter des anticorps dirigés contre le virus de l'hépatite C (HCV) dans un constituant corporel de mammifère suspecté de contenir lesdits anticorps comprenant la mise en contact dudit constituant corporel avec une combinaison d'antigènes de HCV dans des conditions qui permettent les réactions anticorps -
55 antigènes et la détection de la présence de complexes immuns desdits anticorps et desdits antigènes, dans laquelle les antigènes sont dans un ou plusieurs polypeptides obtenus par synthèse chimique ou par expression recombinante, immobilisés à la surface d'une matrice solide convenant pour la détection de HCV par un test immunologique, comprenant :

- (a) un premier antigène de HCV comprenant un épitope du domaine C de la polyprotéine de HCV ; et
- (b) un second antigène de HCV choisi parmi

- (i) un antigène de HCV comprenant un épitope du domaine S ;
- (ii) un antigène de HCV comprenant un épitope du domaine NS3 ;
- (iii) un antigène de HCV comprenant un épitope du domaine NS4 ; et
- (iv) un antigène de HCV comprenant un épitope du domaine NS5 ;

à la condition que la combinaison ne soit pas le peptide p1 avec C100-3, le peptide p35 avec C100-3, le peptide p99 avec C100-3 ou un peptide ayant les acides aminés 9 à 177 de la polyprotéine de HCV-1.

2. Méthode selon la revendication 1 dans laquelle ledit second antigène de HCV comprend un épitope du domaine S et laquelle combinaison comprend en outre au moins un antigène supplémentaire de HCV choisi parmi :

- (i) un antigène de HCV comprenant un épitope du domaine NS3 ;
- (ii) un antigène de HCV comprenant un épitope du domaine NS4 ; et
- (iii) un antigène de HCV comprenant un épitope du domaine NS5 ;

3. Méthode selon la revendication 2 dans laquelle l'antigène supplémentaire provient du domaine NS3.

4. Méthode selon la revendication 3 dans laquelle le premier antigène de HCV est C22 et l'antigène supplémentaire de HCV est C33c.

5. Méthode selon la revendication 2 dans laquelle l'antigène supplémentaire provient du domaine NS4.

6. Méthode selon la revendication 5 dans laquelle le premier antigène de HCV est C22 et l'antigène supplémentaire de HCV est C100.

7. Méthode selon l'une quelconque des revendications 1 à 6 dans laquelle l'antigène du domaine S est l'antigène S2.

8. Méthode selon l'une quelconque des revendications 1 à 7 dans laquelle la combinaison est sous la forme d'un polypeptide de fusion.

9. Méthode selon l'une quelconque des revendications 1 à 8 dans laquelle la combinaison est sous la forme dudit premier antigène de HCV et lesdits antigènes supplémentaires sont liés à une matrice solide commune.

10. Méthode selon la revendication 9 dans laquelle la matrice solide est la surface d'un puits d'une plaque de micro-titration, une bille ou une bandelette.

11. Combinaison d'antigènes du virus de l'hépatite C (HCV) en un ou plusieurs polypeptides obtenus par synthèse chimique ou expression recombinante, comprenant :

- (a) un premier antigène de HCV comprenant un épitope du domaine C de la polyprotéine de HCV ; et
- (b) un second antigène de HCV choisi parmi

- (i) un antigène de HCV comprenant un épitope du domaine S ;
- (ii) un antigène de HCV comprenant un épitope du domaine NS3 ;
- (iii) un antigène de HCV comprenant un épitope du domaine NS4 ; et
- (iv) un antigène de HCV comprenant un épitope du domaine NS5 ;

à la condition que la combinaison ne soit pas le peptide p1 avec C100-3, le peptide p35 avec C100-3, le peptide p99 avec C100-3 ou un peptide ayant les acides aminés 9 à 177 de la polyprotéine de HCV 1.

12. Combinaison selon la revendication 12 dans laquelle ledit second antigène de HCV comprend un épitope du domaine S et laquelle combinaison comprend en outre au moins un antigène de HCV supplémentaire choisi parmi :

- (i) un antigène de HCV comprenant un épitope du domaine NS3 ;
- (ii) un antigène de HCV comprenant un épitope du domaine NS4 ; et

(iii) un antigène de HCV comprenant un épitope du domaine NS5 ;

13. Combinaison selon la revendication 12 dans laquelle l'antigène supplémentaire provient du domaine NS3.

5 14. Combinaison selon la revendication 13 dans laquelle le premier antigène de HCV est C22 et l'antigène de HCV supplémentaire est C33c.

15. Combinaison selon la revendication 14 dans laquelle l'antigène supplémentaire provient du domaine NS4.

10 16. Combinaison selon la revendication 15 dans laquelle le premier antigène de HCV est C22 et l'antigène supplémentaire de HCV est C100.

17. Combinaison selon l'une quelconque des revendications 11 à 16 dans laquelle l'antigène du domaine S est l'antigène S2.

15 18. Combinaison selon l'une quelconque des revendications 11 à 17 dans laquelle la combinaison est sous la forme d'un polypeptide de fusion.

20 19. Combinaison selon l'une quelconque des revendications 11 à 18 dans laquelle la combinaison est sous la forme dudit premier antigène de HCV et lesdits antigènes supplémentaires sont liés à une matrice solide commune.

20. Combinaison selon la revendication 19 dans laquelle la matrice solide est la surface d'un puits d'une plaque de microtitration, un bille ou une bandelette.

25 21. Combinaison selon l'une quelconque des revendications 11 à 20 dans laquelle la combinaison est immobilisée à la surface d'une matrice convenant pour la détection de HCV par un test immunologique.

30 22. Kit pour réaliser un test de détection des anticorps dirigés contre un antigène de l'hépatite C (HCV) dans un constituant corporel de mammifère suspecté de contenir lesdits comprenant, sous forme de combinaison conditionnée :

(a) la combinaison selon l'une quelconque des revendications 11 à 21 ;

(b) les réactifs témoins standard ; et

(c) les instructions pour réaliser le test.

35

40

45

50

55

FIG. 1A

-341 GCCAGCCCCCTGATGGGGCGA
CGGTCGGGGGACTACCCCCGCT

-319 CACTCCACCATGAATCACTCCCCCTGTGAGGAACTACTGTCTTACGCAGAAAGCGTCTAG
GTGAGGTGGTACTTAGTGAGGGGACACTCCTTGATGACAGAAAGTGCGTCTTTCGCAGATC

-259 CCATGGCGTTAGTATGAGTGTCTGTCAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATA
GGTACCGCAATCATCTACAGCACGTCGGAGGTCTTGGGGGGAGGGCCCTCTCGGTAT

-199 GTGGTCTGCGGAAACCGGTGAGTACACCGGAATTGCCAGGACGACGGGTCTTCTTGTGA
CACCAGACGCTTGGCCACTCATGTGGCCCTTAACGGTCTGTGCTGCCAGGAAAGAACCT

-139 TCAACCCGCTCAATGCCCTGGAGATTGGGCGTGCCCCCGCAAGACTGTAGCCCGAGTAGT
AGTTGGGCGAGTTACGACCTCTAAACCCGCACGGGGCGTTCTGACGATCGGCTCATCA

- 79 GTTGGGTGCGGAAAGGCCTTGTGGTACTGCCCTGATAGGGTGCTTGCAGTCCCCGGGAG
CAACCCAGCGCTTCCGGAACACCATGACGGACTATCCACGAAACGCTCACGGGGCCCTC

- 19 GTCTCGTAGACCGTGCACC
CAGAGCATCTGGCACGTGG

Arg Thr

MetSerThrAsnProLysProLysProGlnLysLysAsnLysArgAsnThrAsnArgArgProGln
1 ATGAGCACGAATCCCTAAACCTCAAAAAAACAACGTAACACCAACCGTCGCCCCACAG
TACTCGTGCTTAGGATTGGAGTTTTTTTTTTGTGTCATTGTGGTTGGCAGCGGGTGTC

AspValLysPheProGlyGlyGlyGlnIleValGlyGlyValTyrLeuLeuProArgArg
61 GACGTCAAAGTTCGGGTGGCGGTGAGATCGTTGGTGGAGTTTACTTGTTCGCCGCAGG
CTGCAGTTC AAGGGCCACCGCCAGTCTAGCAACCCACCTCAAATGAACAACGGCGGCTCC

GlyProArgLeuGlyValArgAlaThrArgLysThrSerGluArgSerGlnProArgGly
 121 GGCCCTAGATTGGGTGTGGCGCGACGAGAAAGACTTCCGAGCGGTCCGAACCTCGAGGT
 CCGGGATCTAACCCACACGCGCGCTGCTCTTTCTGAAGGCTCGCCAGCGTTGGAGCTCCA

 ArgArgGlnProIleProLysAlaArgArgProGluGlyArgThrTrpAlaGlnProGly
 181 AGACGTCAGCCTATCCCCAAGGCTCGTCGGCCCGAGGGCAGGACCTGGGCTCAGCCCCGG
 TCTGCAGTCGGATAGGGTTCCGAGCAGCCGGGCTCCCGTCTCTGGACCCGAGTCGGGCCC

 TyrProTrpProLeuTyrGlyAsnGluGlyCysGlyTrpAlaGlyTrpLeuLeuSerPro
 241 TACCCCTTGGCCCTCTATGGCAATGAGGGCTGCGGGTGGCGGATGGCTCCTGTCTCCC
 ATGGGAACCGGGAGATACCGTTACTCCCGACGCCACCCGCCCTACCGAGGACAGAGGG

 ArgGlySerArgProSerTrpGlyProThrAspProArgArgSerArgAsnLeuGly
 301 CGTGGCTCTCGGCCCTAGCTGGGCCCCACAGACCCCGCGTAGGTCCGCAATTGGGT
 GCACCGAGAGCCGGATCGACCCCGGGTGTCTGGGGCCGCATCCAGCGGTTAAACCCA

 LysValIleAspThrLeuThrCysGlyPheAlaAspLeuMetGlyTyrIleProLeuVal
 361 AAGGTCATCGATACCCCTTACGTGGGCTTCGCCGACCTCATGGGGTACATACCGCTCGTC
 TTCCAGTAGCTATGGGAATGCACGCCCGAAGCGGCTGGAGTACCCCATGTATGGCGAGCAG

 GlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAlaHisGlyValArgValLeuGluAsp
 421 GCGCCCCCTCTTGGAGGCGCTGCCAGGCCCTGGCGCATGGCGTCCGGGTTCTGGAAGAC
 CCGCGGGGAGAACCTCCGCGACGGTCCCGGACCGCGTACCGCAGGCCCAAGACCTTCTG

FIG. 1B

FIG. 1C

Thr

481 GlyValAsnTyrAlaThrGlyAsnLeuProGlyCysSerPheSerIlePheLeuLeuAla
 GCGTGAACTATGCAACAGGGAACCTTCCTGGTTGCTCTTCTCTATCTTCTTCTGGCC
 CCGCACTTGATACTGTTCTCCCTTGAAGGACCAACGAGATGATAGAGGAAGACCGG

541 LeuLeuSerCysLeuThrValProAlaSerAlaTyrGlnValArgAsnSerThrGlyLeu
 CTGCTCTCTTGCTTGACTGTGCCCCGCTTCGGCCTACCAAGTGCGCAACTCCACGGGGCTT
 GACGAGAGAACGAACCTGACACGGGGGAGCGGATGGTTACGCGTTGAGGTGCCCGGAA

601 TyrHisValThrAsnAspCysProAsnSerSerIleValTyrGluAlaAlaAspAlaIle
 TACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGGCGCGATGCCATC
 ATGGTGCACTGGTTACTAACGGGATTGAGCTCATAACACATGCTCCGCCGCTACGGTAG

661 LeuHisThrProGlyCysValProCysValArgGluGlyAsnAlaSerArgCysTrpVal
 CTGCACACTCCGGGTGCGTCCCTTGCCTTCGTGAGGGCAACGCCCTCGAGGTGTTGGGTG
 GACGTGTGAGGCCCCACGCAGGGAACGCAAGCACTCCCGTTGCGGAGCTCCACAACCCAC

721 AlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThrGlnLeuArgArg
 GCGATGACCCCTACGGTGGCCACCAGGATGGCAAACTCCCGCGACGACGCTTCGACGT
 CGCTACTGGGGATGCCACCGGTGGTCCCTACCGTTTGAGGGGCGCTGCGTGAAGCTGCA

781 HisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyrValGlyAspLeu
 CACATCGATCTGCTTGTCGGGAGCGCCACCCTCTGTTGCGGCCCTCTACGTGGGGACCTA
 GTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGGGAGATGCACCCCTGGAT

841 CysGlySerValPheLeuValGlyGlnLeuPheThrPheSerProArgArgHisTrpThr
 TGCGGGTCTGTCTTCTTGTCGGCCAACTGTTCACCTTCTCTCCAGGCGCCACTGGACG
 ACGCCCAGACAGAAAGAACAGCCCGGTTGACAAAGTGGAAGAGAGGGTCCCGGTGACCTGC

901 ThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrp
 ACGCAAGGTTGCAATTGCTCTATCTATCCCGGCATATAACGGGTACCGCATGGCATGG
 TCGGTTCCAACGTTAACGAGATAGATAGGGCCGTATATTGCCAGTGGCGTACCGTACC

 961 AspMetMetMetAsnTrpSerProThrAlaLeuValMetAlaGlnLeuLeuArgIle
 GATATGATGATGAACCTGGTCCCTACGACGGCGTTGGTAATGGCTCAGCTGCTCCGGATC
 CTATACTACTACTTGACACGAGGATGCTGCCGCAACCATTAACGAGTCGACGAGGCCCTAG
 Val
 1021 ProGlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAla
 CCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTGGGGAGTCCCTGGCGGCATAGCG
 GGTGTTCCGGTAGAACCTGTACTAGCACGACGAGTGACCCCTCAGGACCGCCCGTATCGC
 1081 TyrPheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGly
 TATTTCTCCATGGTGGGAACTGGGCGAAGTCTCTGGTAGTGCTGCTGCTATTGCGCGC
 ATAAAGAGGTACCAACCCCTTGACCCGCTTCCAGGACCATCACGACGACGATAAACGGCCG
 1141 ValAspAlaGluThrHisValThrGlyGlySerAlaGlyHisThrValSerGlyPheVal
 GTCGACGCGGAAACCCACGTCACCGGGGAAAGTCCCGGCCACACTGTGTCTGGATTGTT
 CAGCTGCGCCTTTGGGTGCAGTGGCCCCCTTCACGGCCGGTGTGACACAGACCTAAACAA
 1201 SerLeuLeuAlaProGlyAlaLysGlnAsnValGlnLeuIleAsnThrAsnGlySerTrp
 AGCCTCCTCGCACCGCGGCCAAGCAGAACGTCCAGCTGATCAACACCAACGGCAGTTGG
 TCGGAGGAGCGTGGTCCCGCGTTCTGCTGTGAGGTCGACTAGTTGTGTGTCCTCAACC

FIG. 1D

FIG. 1E

1261 HisLeuAsnSerThrAlaLeuAsnCysAsnAspSerLeuAsnThrGlyTrpLeuAlaGly
 CACCTCAATAGCAGGCCCTGAACCTGCAATGATAGCCTCAACACCGGCTGGTTGGCAGGG
 GTGGAGTTATCGTGCCGGGACTTGACGTTACTATCGGAGTTGTGGCCGACCAACCGTCCC

1321 LeuPheTyrHisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArg
 CTTTTCATATCACCAAGTTCAACTCTTCAGGCTGTCTGAGAGGCTAGCCAGCTGCCGA
 GAAAAGATAGTGGTGTTCAGTTGAGAACTCCGACAGGACTCTCCGATCGGTCGACGGCT

1381 ProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyPro
 CCCCTTACCGATTTTGACCAAGGCTGGGGCCCTATCAGTTATGCCAACGGAAGCGCCCC
 GGGGAATGGCTAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCCTTCGCCGGGG

1441 AspGlnArgProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLys
 GACCAGCGCCCTACTGCTGGCACTACCCCCCAAACCTTGCGGTATTGTGCCCGCGAAG
 CTGGTCGCGGGGATGACGACCGTGATGGGGGTTTGTGGAACGCCATAAACACGGGCGCTTC

1501 SerValCysGlyProValTyrCysPheThrProSerProValValGlyThrThrAsp
 AGTGTGTGTGGTCCGGTATATTGCTTCACTCCAGCCCGTGGTGGTGGAAACGACCGAC
 TCACACACACACAGGCCATATAACGAAGTGAGGGTCGGGGCACCAACCCTTGCTGGCTG

1561 ArgSerGlyAlaProThrTyrSerTrpGlyGluAsnAspThrAspValPheValLeuAsn
 AGGTCGGGCGCGCCACCTACAGCTGGGGTGAAATGATACGGACGCTCTCGTCCCTAAC
 TCCAGCCCCGCGGGTGATGTCGACCCCCACTTTTACTATGCTGCAGAAAGCAGGAATTG

1621 AsnThrArgProProLeuGlyAsnTrpPheGlyCysThrTrpMetAsnSerThrGlyPhe
 AATACCAGGCCACCGCTGGCAATTGGTTCGGTTGTACCTGGATGAACCTCAACTGGATTTC
 TTATGTGTCGGTGGCGACCCGTTAAACCAAGCCAAACATGGACCTACTTGAGTTGACCTAAG

1681 ThrLysValCysGlyAlaProProCysValIleGlyGlyAlaGlyAsnAsnThrLeuHis
 ACCAAAGTGTGGGAGCGCCTCTTGTGTATCGGAGGGGGGCAACAAACACCTGCAC
 TGGTTTACACGCCCTCGCGGAGGAACACAGTAGCCTCCCCCGCTTGTGTGGGACGTG

 1741 CysProThrAspCysPheArgLysHisProAspAlaThrTyrSerArgCysGlySerGly
 TGCCCCACTGATTGCTTCCGCAAGCATCCGGACGCCACATACCTCTCGGTGGCTCCGGT
 ACGGGTGACTAAACGAAGCGTTCGTAGGCCCTGGGTGTATGAGAGCCACGCCGAGGCCA

 Ile
 1801 ProTrpLeuThrProArgCysLeuValAspTyrProTyrArgLeuTrpHisTyrProCys
 CCTGGATCACACCCAGGTGCCCTGGTCGACTACCCGTATAGGCTTTGGCATATACCTTGT
 GGGACCTAGTGTGGGTCCACGGACCAGCTGATGGGCATATCCGAAACCGTAATAGGAACA

 1861 ThrIleAsnTyrThrIlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeu
 ACCATCAACTACACCATATTTAAATCAGGATGTACGTGGAGGGGTCCGAACACACAGGCTG
 TGGTAGTTGATGTGGTATAAAATTTTAGTCCTACATGCACCCCTCCCCAGCTTGTGTCCGAC

 1921 GluAlaAlaCysAsnTrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSer
 GAAGCTGCCCTGCAACTGGACGCGGGCGAACCCTTGCATCTGGAAGACAGGGACAGGTCC
 CTTCCGACGGACGTTGACCTGCGCCCCCGCTTGCAACGCTAGACCTTCTGTCCCTGTCCAGG

 1981 GluLeuSerProLeuLeuLeuThrThrThrGlnTrpGlnValLeuProCysSerPheThr
 GAGCTCAGCCCGTTACTGCTGACCACTACACAGTGGCAGGTCTCCCGTGTTCCTTCACA
 CTCGAGTCGGGCAATGACGACTGGTGTGTGTACCGTCCAGGAGGGCACAAAGGAAGTGT

FIG. 1F

FIG. 1G

2041 ThrLeuProAlaLeuSerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGln
 ACCCTACCAGCCTTGTCACCGGCTCATCCACCTCCACAGAACATTTGTGACGTGCAG
 TGGGATGGTCGGAACAGGTGGCCGAGTAGTGGAGTGCTTGTAAACACCTGCACGTC

 2101 TyrLeuTyrGlyValGlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValVal
 TACTTGTAACGGGTGGGTCAAGCATCGCGTCTCGTCCCTGGGCCATTAAAGTGGAGTACGTCGTT
 ATGAACATGCCCCACCCAGTTCTGTAGCGCAGGACCCGGTAATTCAACCCTCATGCAGCAA

 2161 LeuLeuPheLeuLeuAlaAspAlaArgValCysSerCysLeuTrpMetMetLeuLeu
 CTCCTGTTCCCTTGCTGTCAGACGCGCGCTCTGCTCCTGCTTGTGGATGATGCTACTC
 GAGGACAAAGGAAGACGAAACGTCTGCGCGCGCAGACGAGGACGAACACCTACTACGATGAG

 2221 IleSerGlnAlaGluAlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAla
 ATATCCCAAGCGAGCGGCTTTGGAGAACCTCGTAATACTTAATGCAGCATCCCTGGCC
 TATAGGGTTCCGCTCCGCCGAAACCTCTTGGAGCATTATGAATTACGTCGTAGGGACCGG

 2281 GlyThrHisGlyLeuValSerPheLeuValPhePheCysPheAlaTrpTyrLeuLysGly
 GGGACGCACGGTCTTGATATCCTTCCTCGTGTCTTCTGCTTTGCCATGGTATTTGAAGGGT
 CCCTGCGTGCCAGAACATAGGAAGGAGCACAAAGAACGAAACGTACCATAACTTCCCA

 2341 LysTrpValProGlyAlaValTyrThrPheTyrGlyMetTrpProLeuLeuLeuLeu
 AAGTGGGTGCCCCGAGCGGTCTACACCTTCTACGGGATGTGGCCCTCTCCTGCTCCTG
 TTCACCCACGGGCTCGCCAGATGTGGAAGATGCCCTACACCGGAGAGGACGAGGAC

 2401 LeuAlaLeuProGlnArgAlaTyrAlaLeuAspThrGluValAlaAlaSerCysGlyGly
 TTGGCGTTGCCCCAGCGGCGTACCGCTGGACACGAGGTGGCCGCTCGTGTGGCGGT
 AACCGCAACGGGGTCCGCCCGCATGCGCGACCTGTGCTCTCCACCGGCGCAGCACACCGCCA

2461 ValValLeuValGlyLeuMetAlaLeuThrLeuSerProTyrTyrLysArgTyrIleSer
 GTTGTTCCTCGTGGGTTGATGGCGCTGACTCTGTCAACCATATTAACAAGCGCTATATCAGC
 CAACAAGAGCAGCCCAACTACCGGACTGAGACAGTGGTATAATGTTCCGGATATAGTCG

 (Asn)
 2521 TrpCysLeuTrpTrpLeuGlnTyrPheLeuThrArgValGluAlaGlnLeuHisValTrp
 TGGTGCTTGTGGCTTCAGTATTTTCTGACCAAGAGTGGAAGCGCAACTGCACGTGTGG
 ACCACGAACACACCGAAGTCATAAAGACTGGTCTCACCTTCGCGTTGACGTGCACACC

 2581 IleProProLeuAsnValArgGlyGlyArgAspAlaValIleLeuLeuMetCysAlaVal
 ATTCCCCCCTCAACGTCGAGGGGCGGACGCCGTCACTTACTCATGTGTGCTGTA
 TAAGGGGGGAGTTGCAGGCTCCCCCGCTGCGGAGTAGAATGAGTACACACGACAT

 2641 HisProThrLeuValPheAspIleThrLysLeuLeuAlaValPheGlyProLeuTrp
 CACCCGACTCTGGTATTTGACATCACCAAAATTGCTGCTGGCGCTTTCGACCCCTTGG
 GTGGGCTGAGACCATAAACTGTAGTGGTTAACGACGACCGGCAGAACCTGGGGAAACC

 2701 IleLeuGlnAlaSerLeuLeuLysValProTyrPheValArgValGlnGlyLeuLeuArg
 ATTCTTCAAGCCAGTTTGCTTAAAGTACCCTACTTGTGCGCGTCCAAGGCCCTCTCCGG
 TAAGAAAGTTCGGTCAAACGAAATTTTCATGGGATGAACACGCGCAGGTTCCGGGAAGAGGCC

 2761 PheCysAlaLeuAlaArgLysMetIleGlyGlyHisTyrValGlnMetValIleIleLys
 TTCTGCGCGTTAGCGGGAAGATGATCGGAGGCCATTACGTGCAAAATGGTCATCATTAAG
 AAGACGCGCAATCGCGCCTTCTACTAGCCTCCGGTAATGCACGTTTACCAGTAGTAATTC

FIG. 1H

FIG. 11

2821	LeuGlyAlaLeuThrGlyThrTyrValTyrAsnHisLeuThrProLeuArgAspTrpAla TTAGGGCGCTTACTGGCACCTATGTTTATAAACCATCTCACTCTCTTCGGGACTGGCGG AATCCCCGGAATGACCGTGATACAAATATTTGGTAGAGTGAGGAGAACCCCTGACCCCGC
2881	HisAsnGlyLeuArgAspLeuAlaValAluProValValPheSerGlnMetGlu CACAAACGGCTTGCGAGATCTGGCCGTGGCTGTAGAGCCAGTCGTCTTCTCCCAAATGGAG GTGTTGCCGAACGCTCTAGACCGGCACCGACATCTCGGTACGACAGAGGGTTTACCTC
2941	ThrLysLeuIleThrTrpGlyAlaAspThrAlaAlaCysGlyAspIleIleAsnGlyLeu ACCAAGCTCATCACGTGGGGGCGAGATACCGCCGCGTGGTGACATCATCAACGGCTTG TGGTTCGAGTAGTGCACCCCGCTCTATGGGGCGCACGCCACTGTAGTAGTTGCCGAAC
3001	ProValSerAlaArgGlyArgGluIleLeuLeuGlyProAlaAspGlyMetValSer CCTGTTTCCCGCCGCGAGGGCGGAGATACTGCTCGGGCCAGCCGATGGAATGGTCTCC GGACAAAGCGGGCGTCCCCGGCCCTCTATGACGAGCCCGGTCGGCTACCTTACCAGAGG
3061	LysGlyTrpArgLeuAlaProIleThrAlaTyrAlaGlnGlnThrArgGlyLeuLeu AAGGGTGGAGGTGCTGGCGCCCATCACGGCGTACGCCCCAGCACACAAGGGCCCTCCTA TTCCCCACCTCCAAACGACCGCGGTAGTCCCGCATGCGGGTCTGTCTTCCCCGGAGGAT
3121	GlyCysIleIleThrSerLeuThrGlyArgAspLysAsnGlnValGluGlyGluValGln GGGTGCATAATCACACGCTAACTGGCCGGACAAACCAAGTGAGGGTGAGGTCCAG CCCACGTATTAGTGGTCCGATTGACCGGCCCTGTTTGGTTCACCTCCCACTCCAGGTC
3181	IleValSerThrAlaAlaGlnThrPheLeuAlaThrCysIleAsnGlyValCysTrpThr ATTGTGTCAACTGCTGCCCAACCTTCTCTGGCAACGTGCATCAATGGGTGTGCTGGACT TAACACAGTTGACGACGGGTTTGGAAAGGACCGTTGCACGTAGTTACCCACACGACCTGA

3241 ValTyrHisGlyAlaGlyThrArgThrIleAlaSerProLysGlyProValIleGlnMet
 GTCTACCAACGGGCGGAAACGAGGACCATCGCGTCACCCAAAGGTCCCTGTCATCCAGATG
 CAGATGGTGCCCCCGCCTTGCTCTCTGCTAGCGCAGTGGGTTCGCCAGGACAGTAGGCTCTAC

 3301 TyrThrAsnValAspGlnAspLeuValGlyTrpProAlaProGlnGlySerArgSerLeu
 TATACCAATGTAGACCAAGACCTTGTGGCTGGCCCGCTCCGCAAGGTAGCCGCTCATTTG
 ATATGGTTACATCTGTTCTGGAACACCCGACCGGGCGAGGCTTCCATCGGGCGAGTAAC

 3361 ThrProCysThrCysGlySerSerAspLeuTyrLeuValThrArgHisAlaAspValIle
 ACACCCCTGCACCTTGGGCTCCTCGGACCTTTACCTGGTCACGAGGCACGCCGATGTCAATT
 TGTGGGACGTGAACGCCCGAGGAGCCTGGAAATGGACCAAGTCCGTCCGTGGGCTACAGTAA

 3421 ProValArgArgGlyAspSerArgGlySerLeuLeuSerProArgProIleSerTyr
 CCGGTGCGCGCGGGGTGATAGCAGGGGCAGCCCTGCTGTCGCCCCCGCCCATTTCCCTAC
 GGGCACGGCGCGCCCCACTATCGTCCCCCGTCGGACGACAGCGGGCGCGGTAAAGGATG

 3481 LeuLysGlySerSerGlyGlyProLeuLeuCysProAlaGlyHisAlaValGlyIlePhe
 TTGAAAGGCTCCTCGGGGGTCCGCTGTTGTGCCCCCGGGGCACGCCGTGGCATATTT
 AACTTCCGAGGAGCCCCCAGGCGAACACAGGGGCGCCCCGTGGGCACCCCGTATAAA

 3541 ArgAlaAlaValCysThrArgGlyValAlaLysAlaValAspPheIleProValGluAsn
 AGGGCCGGGTGTGCACCCGTGGAGTGGCTAAGCGGTGACTTTATCCCTGTGGAGAAC
 TCCCGGCGCCACACGTGGGCACCTCACCGATTCCGCCACCTGAAATAGGGACACCTCTTG

FIG. 1J

FIG. 1K

3601 LeuGluThrThrMetArgSerProValPheThrAspAsnSerSerProProValValPro
 CTAGAGACAACCATGAGTCCCCGGTGTTCACGGATAAATCTCTCCACCATGAGTCCCC
 GATCTCTGTTGGTACTCCAGGGGCCAACAGTGCCTATTGAGGAGAGGTGTCATCACGGG

 3661 GlnSerPheGlnValAlaHisLeuHisAlaProThrGlySerGlyLysSerThrLysVal
 CAGAGCTTCCAGGTGGCTCACCTCCATGCTCCACAGGCGCGCAAAAGCACCAAGGTC
 GTCTCGAAGTCCACCGAGTGGAGGTACGAGGGTGTCGTCGCCGTTTTCGTGGTTCCAG

 3721 ProAlaAlaTyrAlaAlaGlnGlyTyrLysValLeuValLeuAsnProSerValAlaAla
 CCGGCTGCATATGCAGCTCAGGGCTATAAGGTGCTAGTACTCAACCCCTCTGTGCTGCA
 GCGGACGTATACGTGAGTCCCGATATTTCCACGATCATGAGTTGGGGAGACAAACGACGT

 3781 ThrLeuGlyPheGlyAlaTyrMetSerLysAlaHisGlyIleAspProAsnIleArgThr
 AACTGGGCTTTGGTGCTTACATGTCCAAGGCTCATGGGATCGATCCTAACATCAGGACC
 TGTGACCCGAAACCAACGAATGTACAGGTTCCGAGTACCCCTAGCTAGGATTGTAGTCTTGG

 3841 GlyValArgThrIleThrThrGlySerProIleThrTyrSerThrTyrGlyLysPheLeu
 GGGTGAGAACAAATTACCACTGGCAGCCCCCATCAGTACTCCACCTACGGCAAGTTCCTT
 CCCCACCTCTGTGTTAATGTTGACCGTCCGGGTAGTGATGAGGTGGATGCCGTTCAAGGAA

 3901 AlaAspGlyGlyCysSerGlyGlyAlaTyrAspIleIleCysAspGluCysHisSer
 GCCGACGGCGGTGCTCGGGGGCGGCTTATGACATAATAATTGTGACGAGTGCCACTCC
 CGGCTGCCGCCCCACGAGCCCCCGGAATACTGTATATTATAAACACTGCTCACGGTGAGG

(Val)
 ThrAspAlaThrSerIleLeuGlyIleGlyThrValLeuAspGlnAlaGluThrAlaGly
 3961 ACGGATGCCACATCCATCTTGGGCATCGGCACCTGTCTTACCAAGCAGAGACTGCGGGG
 TGCCTACGGGTAGGTAGAACCCGTAGCCGTGACAGGAACCTGGTTCGTCTCTGACGCCCC
 AlaArgLeuValValLeuAlaThrAlaThrProProGlySerValThrValProHisPro
 4021 GCGAGACTGGTTGTGCTCGCCACCGCCACCCCTCCGGGCTCCGTCACTGTGCCCATCCC
 CGCTCTGACCAACACGAGCGGTGGGTGGGAGGCCCGAGGCAGTGACACGGGGTAGGG
 AsnIleGluGluValAlaLeuSerThrThrGlyGluIleProPheTyrGlyLysAlaIle
 4081 AACATCGAGGAGTTGTCTGTCCACCCAGGAGAGATCCCTTTTACGGCAAGGCTATC
 TTGTAGCTCCTCCAACGAGACAGGTGGTGGCCTCTCTAGGAAATAATGCCGTTCCGATAG
 ProLeuGluValIleLysGlyGlyArgHisLeuIlePheCysHisSerLysLysCys
 4141 CCCCTCGAAGTAATCAAGGGGGGAGACATCTCATCTTCTGTCTTCAATCAAGAAGATGC
 GGGAGCTTCATTAGTTCCCCCTCTGTAGAGTAGAAGACAGTAAGTTCTTCTTCACG
 AspGluLeuAlaAlaLysLeuValAlaLeuGlyIleAsnAlaValAlaTyrTyrArgGly
 4201 GACGAACTCGCCGCAAGCTGGTCGCATTGGGCATCAATGCCGTGGCCTACTACCGCGGT
 CTGCTTGAGCGGCGTTTCGACCCAGCGTAACCCGTAGTTACGGCACCGGATGATGGGCCCA
 LeuAspValSerValIleProThrSerGlyAspValValValAlaThrAspAlaLeu
 4261 CTTGACGTGTCCGTCAATCCGACCCAGCGGCGATGTTGTCTGTCTGGCAACCGATGCCCTC
 GAACTGCACAGGCAGTAGGGCTGGTCCGCCCTACAACAGCAGCACCGTTGGCTACGGGAG
 Tyr
 MetThrGlyTyrThrGlyAspPheAspSerValIleAspCysAsnThrCysValThrGln
 4321 ATGACCGGCTATACCGGCGACTTCGACTCGGTGATAGACTGCAATACGTGTCAACCCAG
 TACTGGCCGATATGGCCGCTGAAGCTGAGCCACTATCTGACGTTATGCACACAGTGGGTC

FIG. 1L

FIG. 1M

(Ser)

ThrValAspPheSerLeuAspProThrPheThrIleGluThrIleThrLeuProGlnAsp
 4381 ACAGTCGATTTCAGCCCTTGACCCCTACCTTCACCATTGAGACAAATCACGCTCCCCCAGGAT
 TGTCAGCTAAAGTCGGAACTGGGATGGAAAGTGTAACCTCTGTAGTGCAGGGGGTCCCTA

 AlaValSerArgThrGlnArgArgGlyArgThrGlyArgGlyLysProGlyIleTyrArg
 4441 GCTGTCTCCCGCACTCAACGTCGGGGCAGGACTGGCAGGGGAAAGCCAGGCATCTACAGA
 CGACAGAGGGCGTGAGTTGCAGCCCCCGTCCCTGACCGTCCCCCTTCGGTCCGTAGATGTCT

 PheValAlaProGlyGluArgProSerGlyMetPheAspSerSerValLeuCysGluCys
 4501 TTTGTGGCACCGGGGAGCGCCCTCCGGCATGTTCGACTCGTCCGTCTCTGTGAGTGC
 AAACACCGTGGCCCCCTCGCGGGAGGCCGTACAAGCTGAGCAGGACGAGACACTCACG

 TyrAspAlaGlyCysAlaTrpTyrGluLeuThrProAlaGluThrThrValArgLeuArg
 4561 TATGACGACGGCTGTGCTTGGTATGAGCTCACGCCCCCGGAGACTACAGTTAGGCTACGA
 ATACTGCGTCCGACACGAACCATACTCGAGTGGGGCGGCTCTGATGTCAATCCGATGCT

 AlaTyrMetAsnThrProGlyLeuProValCysGlnAspHisLeuGluPheTrpGluGly
 4621 GCGTACATGAACACCCCGGGCTTCCCGTGTGCCAGGACCATCTTGAATTTTGGGAGGGC
 CGCATGTACTTGTGGGGCCCCGAAAGGCACACGCTCCTGTAGAACTTAAACCCCTCCCG

 ValPheThrGlyLeuThrHisIleAspAlaHisPheLeuSerGlnThrLysGlnSerGly
 4681 GTCTTTACAGGCCTCACTCATATAGATGCCCACTTTCTATCCCAGACAAAGCAGAGTGGG
 CAGAAATGTCCGGAGTGAGTATATCTACGGGTGAAAGATAGGGTCTGTCTTCTCACC

 GluAsnLeuProTyrLeuValAlaTyrGlnAlaThrValCysAlaArgAlaGlnAlaPro
 4741 GAGAACCTTCCCTTACCTGGTAGCGTAGCAAGCCACCGTGTGGCTAGGGCTCAAGCCCCCT
 CTCTTGGAAAGGAATGGACCATCGCATGGTTCGGTGGCACACGCCGATCCCGAGTTCGGGGA

4801 ProProSerTrpAspGlnMetTrpLysCysLeuIleArgLeuLysProThrLeuHisGly
 CCCCATCGTGGACCAAGATGTGAAGTGTGATTGCTCGCTCAAGCCACCCCTCCATGGG
 GGGGTAGCACCCCTGGTCTACACCTTCACAACTAAGCGGAGTTCGGGTGGGAGGTACCC

 4861 ProThrProLeuLeuTyrArgLeuGlyAlaValGlnAsnGluIleThrLeuThrHisPro
 CCAACACCCCTGCTATACAGACTGGCGCTGTTTCAGAAATGAATCACCCTGACGCACCCA
 GGTGTGGGACGATATGTCTGACCCCGCACAAAGTCTTACTTTAGTGGGACTGCGTGGGT

 4921 ValThrLysTyrIleMetThrCysMetSerAlaAspLeuGluValValThrSerThrTrp
 GTCACCAATAACATCATGACATGTCATGTCCGCCGACCTGGAGGTCGTACAGCACCTGG
 CAGTGGTTTATGTAGTACTGTACGTACAGCCGGCTGGACCTCCAGCAGTGTCTCGTGACC

 4981 ValLeuValGlyGlyValLeuAlaAlaLeuAlaAlaTyrCysLeuSerThrGlyCysVal
 GTGCTCGTTGGCGGCTCCTGGCTGCTTGGCCCGGTATTGCCCTGTCAACAGGCTGCGTG
 CACGAGCAACCCCGCAGGACCGACGAAACCGCGCATAAACGGACAGTTGTCCGACGCAC

 5041 ValIleValGlyArgValValLeuSerGlyLysProAlaIleIleProAspArgGluVal
 GTCATAGTGGCAGGTCGTCTTGTCCGGGAAGCCGGCAATCATACCTGACAGGGAAGTC
 CAGTATCACCCGTCCTCCAGCAGAACAGGCCCTTCGGCCGTTAGTATGGACTGTCCCTTCAG

 5101 LeuTyrArgGluPheAspGluMetGluGluCysSerGlnHisLeuProTyrIleGluGln
 CTCTACCGAGAGTTCGATGAGATGGAAGAGTGTCTCTCAGCACTTACCGTACATCGAGCAA
 GAGATGGCTCTCAAGCTACTCTACCTTCTCACGAGAGTCGTGAATGGCATGTAGCTCGTT

FIG. 1N

FIG. 10

GlyMetMetLeuAlaGluGlnPheLysGlnLysAlaLeuGlyLeuLeuGlnThrAlaSer
 5161 GGGATGATGCTCGCCGAGCAGTTCAAGCAGAAGCCCTCGGCCCTCCTGCAGACCGGTCC
 CCCTACTACGAGCGGCTCGTCAAGTTCGTCTTCCGGGAGCCGGAGGACGTCTGGCGCAGG

 ArgGlnAlaGluValIleAlaProAlaValGlnThrAsnTrpGlnLysLeuGluThrPhe
 5221 CGTCAGGCAGAGGTTATCGCCCCCTGCTGTCCAGACCAACTGGCAAAACTCGAGACCTTC
 GCAGTCCGTCTCCAATAGCGGGGACGACAGGTCTGTGTGACCGTTTGTGAGCTCTGGAAG

 TrpAlaLysHisMetTrpAsnPheIleSerGlyIleGlnTyrLeuAlaGlyLeuSerThr
 5281 TGGCGGAAGCATATGTGGAACCTTCATCAGTGGGATACAATACTTGGCGGGCTTGTC AACG
 ACCCGCTTCGTATACACCTTGAAGTAGTCAACCTATGTTATGAAACCGCCCCGAACAGTTGC

 LeuProGlyAsnProAlaIleAlaSerLeuMetAlaPheThrAlaAlaValThrSerPro
 5341 CTGCCCTGGTAACCCCGCCATTGCTTCATTGATGGCTTTTACAGCTGCTGTACCCAGCCCA
 GACGGACCATTTGGGGCGGTAAACGAAGTAACTACCCGAAATGTTCGACGACAGTGGTCGGGT

 LeuThrThrSerGlnThrLeuLeuPheAsnIleLeuGlyGlyTrpValAlaAlaGlnLeu
 5401 CTAACCACTAGCCAAACCCCTCCTCTTCAACATATTGGGGGGTGGTGGCTGCCCCAGCTC
 GATTGGTGATCGGTTTGGGAGGAGAAGTTGTATAACCCCCCAACCCACCGGCGGTCCGAG

 AlaAlaProGlyAlaAlaThrAlaPheValGlyAlaGlyLeuAlaGlyAlaAlaIleGly
 5461 GCCGCCCCGGTGCCGCTACTGCCCTTTGTGGCGCTGGCTTAGCTGGCGCCGCCATCCGC
 CGCGGGGGCCACGGCGATGACGGAAACACCCGCGACCGAATCGACCCGGCGGTAGCCG

 SerValGlyLeuGlyLysValLeuIleAspIleLeuAlaGlyTyrGlyAlaGlyValAla
 5521 AGTGTGGACTGGGGAAGTCCCTCATAGACATCCTTGACAGGTATGGCGGGCGGTGGCG
 TCACAACCTGACCCCTTCCAGGAGTATCTGTAGGAACGTCCCATACCGCGCCCGCACCCG

5581 GlyAlaLeuValAlaPheLysIleMetSerGlyGluValProSerThrGluAspLeuVal
 (Gly)
 GGAGCTCTTGTGGCATTCAAGATCATGAGCGGTGAGTCCCTCCACGGAGGACCTGGTC
 CCTCGAGAACACCGTAAGTTCTAGTACTCGCCACTCCAGGGGAGGTGCTCCTGGACCAG

 5641 AsnLeuLeuProAlaIleLeuSerProGlyAlaLeuValValGlyValValCysAlaAla
 AATCTACTGCCCCCATCTCTCGCCCGAGCCCTCGTAGTCGGCGTGTCTGTGCAGCA
 TTAGATGACGGCGGTAGGAGAGCGGGCTCGGGAGCATCAGCCGCCACACACGTCGT

 5701 IleLeuArgArgHisValGlyProGlyGluGlyAlaValGlnTrpMetAsnArgLeuIle
 ATACTGCGCGGCACGTTGGCCCGGCGAGGGGCAGTGCAGTGCATGAACCGGCTGATA
 TATGACGCGCGCGTGCAACCGGGCCCGCTCCCCGTCACTACCTACTTGGCCGACTAT

 5761 AlaPheAlaSerArgGlyAsnHisValSerProThrHisTyrValProGluSerAspAla
 GCCTTCGCCCTCCCGGGGAACCATGTTTCCCCCACGCACTACGTGCCGGAGCGATGCA
 CGGAAGCGGAGGGCCCCCTTGGTACAAAGGGGGTGCGTGATGCACGGCCCTCTCGCTACGT

 5821 (HisCys)
 AlaAlaArgValThrAlaIleLeuSerSerLeuThrValThrGlnLeuLeuArgArgLeu
 GCTGCCCGCGTCACTGCCATACTCAGCAGCCCTCACTGTAAACCCAGCTCCTGAGGCGACTG
 CGACGGGCGCAGTGACGGTATGAGTCGTCGGAGTGACATTTGGGTCGAGGACTCCGCTGAC

 5881 HisGlnTrpIleSerSerGluCysThrThrProCysSerGlySerTrpLeuArgAspIle
 CACCAAGTGGATAAGCTCGGAGTGTACCACTCCATGCTCCGGTTCCTGGCTAAGGACATC
 GTGGTCACCTATTTCGAGCCCTCACATGGTGAGGTACGAGGCCAAGGACCGATTCCCTGTAG

FIG. 1P

FIG. 1Q

5941 TrpAspTrpIleCysGluValLeuSerAspPheLysThrTrpLeuLysAlaLysLeuMet
 TGGGACTGGATATGCGAGGTGTGAGCGACTTTAAGACCTGGCTAAAAGCTAAGCTCATG
 ACCCTGACCTATACGCTCCACAACCTCGCTGAAATTCTGGACCGATTTCGATTCCGAGTAC

 6001 ProGlnLeuProGlyIleProPheValSerCysGlnArgGlyTyrLysGlyValTrpArg
 CCACAGCTGCCTGGGATCCCTTTGTGCTCCTGCCAGCGGGTATAAGGGGTCTGGCGGA
 GGTGTCGACGGACCCCTAGGGAAACACAGGACGGTCCGCGCCCATATTCCCCCAGACCGCT

 6061 (Val)
 GlyAspGlyIleMethIleThrArgCysHisCysGlyAlaGluIleThrGlyHisValLys
 GTGGACGGCATCATGCACACTCGCTGCCACTGTGGAGCTGAGATCACTGGACATGTCAA
 CACTGCCGTAGTACGTGTGAGCGACGGTGACACCTCGACTCTAGTGACCTGTACAGTTT

 6121 AsnGlyThrMetArgIleValGlyProArgThrCysArgAsnMetTrpSerGlyThrPhe
 AACGGGACGATGAGGATCGTCGGTCCTAGGACCTGCAGGAACATGTGGAGTGGACCTTC
 TTGCCCCTGCTACTCCTAGCAGCCAGGATCCTGGACGTCCTTGTTACACCTCACCCCTGGAAG

 6181 ProIleAsnAlaTyrThrThrGlyProCysThrProLeuProAlaProAsnTyrThrPhe
 CCCATTAAATGCCCTACACCAGGGCCCCCTGTACCCCCCTTCTGCGCCGAACTACACGTTT
 GGGTAATTACGGATGTGGTGCCCCGGGACATGGGGGGAAGGACGCGGCTTGATGTGCAAG

 6241 AlaLeuTrpArgValSerAlaGluGluTyrValGluIleArgGlnValGlyAspPheHis
 GCGCTATGGAGGTGTCTGCAGAGGAATAATGTGGAGATAAGGCAGGTGGGGACTTCCAC
 CGCGATACCTCCACACAGACGTCTCCTTATACACCTCTATTCCGTCCACCCCTGAAGGTG

 6301 TyrValThrGlyMetThrThrAspAsnLeuLysCysProCysGlnValProSerProGlu
 TACGTGACGGGTATGACTACTGACAAATCTCAAATGCCCCGTGCCAGGTCCCATCGCCCCGAA
 ATGCACTGCCCATACTGATGACTGTAGAGTTTACGGGACACGGTCCAGGGTAGCGGGCTT

6361 PhePheThrGluLeuAspGlyValArgLeuHisArgPheAlaProProCysLysProLeu
 TTTTTCACAGAAATTGGACGGGTGGCCCTACATAGTTTGGCCCCCTGCAAGCCCTTG
 AAAAGTGTCTTAACCTGCCCCACGCGGATGTATCCAAACGCGGGGACGTTCCGGGAAC

 6421 LeuArgGluGluValSerPheArgValGlyLeuHisGluTyrProValGlySerGlnLeu
 CTGCGGAGGAGGTATCATTCAGAGTAGGACTCCACGAATACCCGGTAGGTGCGCAATTA
 GACGCCCTCCTCCATAGTAAGTCTCATCTGAGGTGCTTATGGGCCATCCCAGCGTTAAT

 6481 ProCysGluProGluProAspValAlaValLeuThrSerMetLeuThrAspProSerHis
 CCTTGGAGCCCCGAACCGGACGTGGCCGTGTGACGTCCATGCTCACTGATCCCTCCCAT
 GGAACTCGGGCTTGGCCTGCACCGGCACAACTGCAGGTACGAGTACTAGGGAGGTA

 6541 IleThrAlaGluAlaAlaGlyArgArgLeuAlaArgGlySerProProSerValAlaSer
 ATAACAGCAGAGCGCGCGGCGAAGGTTGGCGAGGGGATCACCCCTCTGTGGCCAGC
 TATTGTCTCTCCGCGCGCGCGCTTCCAAACCGCTCCCCCTAGTGGGGGAGACACCGGTCG

 6601 SerSerAlaSerGlnLeuSerAlaProSerLeuLysAlaThrCysThrAlaAsnHisAsp
 TCCTCGGCTAGCCAGCTATCCGCTCCATCTCTCAAGGCAACTTGACCCGCTAACCATGAC
 AGGAGCCGATCGGTCGATAGGCGAGGTAGAGAGTCCCGTTGAACGTGGCGATTGGTACTG

 6661 SerProAspAlaGluLeuIleGluAlaAsnLeuLeuTrpArgGlnGluMetGlyGlyAsn
 TCCCCTGATGCTAGCTCATAGAGGCCAACCTCCTATGGAGGCAGGAGATGGCGGCAAC
 AGGGGACTACGACTCGAGTATCTCCGGTTGGAGGATACCTCCGTCCTCTACCCCGCGTTCG

FIG. 1R

FIG. 1S

IleThrArgValGluSerGluAsnLysValValIleLeuAspSerPheAspProLeuVal
 6721 ATCACAGGGTTGAGTCAGAAAAAAGTGGTATTCTGGACTCCTTCGATCCGCTTGTG
 TAGTGGTCCCAACTCAGTCTTTTGTTCACCACTAAGACCTAGGAAGCTAGGCGAACAC

 AlaGluGluAspGluArgGluIleSerValProAlaGluIleLeuArgLysSerArgArg
 6781 GCGGAGGAGGACGAGCGGGAGATCTCCGTACCCGAGAAATCCTGCGGAAGTCTCGGAGA
 CGCCTCCTCCTGCTCGCCCTCTAGAGGCATGGCGCTCTTAGGACGCCCTTCAGAGCCTCT

 PheAlaGlnAlaLeuProValTrpAlaArgProAspTyrAsnProProLeuValGluThr
 6841 TTCGCCCCAGGCCCTGCCCGTTTGGCGCGGCGGACTATAACCCCCCGCTAGTGGAGACG
 AAGCGGTCCGGACGGGCAAAACCCCGCCGCGCTGATATTGGGGGGCGGATCACCTCTGC

 TrpLysLysProAspTyrGluProProValValHisGlyCysProLeuProProLys
 6901 TGGAAAAAGCCCGACTACGAACACCTGTGTCTCATGGCTGTCCGCTCCACCTCCAAAG
 ACCTTTTTCGGGCTGATGCTTGTGGACACACAGGTACCGACAGGCGAAGGTGGAGGTTTC

 SerProProValProProProArgLysLysArgThrValValLeuThrGluSerThrLeu
 6961 TCCCCCTCCTGTGCCCTCCGCCCTCGGAAGAAGCGGACGGTGGTCCCTCACTGAATCAACCCTA
 AGGGGAGGACACGGAGCGCGAGCCCTTCTTCGCCCTGCCACCGAGTGACTTAGTTGGGAT

 (Ser)
 SerThrAlaLeuAlaGluLeuAlaThrArgSerPheGlySerSerSerThrSerGlyIle
 7021 TCTACTGCCCTTGGCCGAGCTCGCCACCAAGAGCTTTGGCAGCTCCTCAACTCCGGCATT
 AGATGACGGAAACCGGCTCGAGCGGTGTCTTCGAAACCGTCGAGGAGTTGAAGGCCGTAA

 ThrGlyAspAsnThrThrThrSerSerGluProAlaProSerGlyCysProProAspSer
 7081 ACGGGCGACAATACGACAACATCCTCTGAGCCCCGCCCTTCTGGCTGCCCCCGACTCC
 TGCCCCGCTGTTATGCTGTTGTAGGAGACTCGGGCGGGGAAGACCGGCGGGGCTGAGG

(PheAla)

7141 AspAlaGluSerTyrSerSerMetProProLeuGluGlyGluProGlyAspProAspLeu
 GACGCTGAGTCCTATTCTCCATGCCCCCCCCCTGGAGGGAGCCTGGGATCCGGATCTT
 CTGCGACTCAGGATAAGGAGGTACGGGGGGGACCTCCCCCTCGGACCCCTAGGCCCTAGAA

7201 SerAspGlySerTrpSerThrValSerSerGluAlaAsnAlaGluAspValValCysCys
 AGCGACGGGTCAATGGTCAACGGTCAGTAGTGAGGCCAACCGCGAGGATGTCGTGTGCTGC
 TCGCTGCCCCAGTACCAGTTGCCAGTCATCACTCCGGTTGGCCCTCTACAGCACACGACG

7261 SerMetSerTyrSerTrpThrGlyAlaLeuValThrProCysAlaAlaGluGluGlnLys
 TCAATGTCTTACTCTTGGACAGGCGCACTCGTCACCCCGTGC CGCGGAAACAGAAA
 AGTTACAGAAATGAGAAACCTGTCCGCGTGAGCAGTGCGGACACGCGGCCCTTCTTGTCTTT

7321 LeuProIleAsnAlaLeuSerAsnSerLeuLeuArgHisHisAsnLeuValTyrSerThr
 CTGCCCCATCAATGCACATAAGCAACTCGTTGCTACGTCAACCAATTTGGTGATATCCACC
 GACGGGTAGTTACGTGATTTCGTTGAGCAACGATGCAGTGGTGTAAACACACATAAGGTGG

7381 ThrSerArgSerAlaCysGlnArgGlnLysLysValThrPheAspArgLeuGlnValLeu
 ACCTCACGCAGTGCTTGCCAAAGGCAGAAAGAAAGTCACATTTGACAGACTGCAGTTCTG
 TGGAGTGGGTCACGAACGGTTTCCGTCTTCTTTCAGTGTAAACTGTCTGACGTTCAAGAC

7441 AspSerHisTyrGlnAspValLeuLysGluValLysAlaAlaSerLysValLysAla
 GACAGCCATTACCGAGCAGTACTCAAGGAGGTTAAAGCAGCGGCTCAAAAGTGAAGGCT
 CTGTCGGTAATGGTCCCTGCATGAGTTCCTCCAATTTTCGTCGCGCGCAGTTTTCACCTCCGA

FIG. 1T

FIG. 1U

7501	(Phe) AsnLeuLeuSerValGluGluAlaCysSerLeuThrProProHisSerAlaLysSerLys AACTTGCTATCCGTAGAGGAAGCTTGCCAGCCTGACGCCCCCACCACACTCAGCCAAATCCAAG TTGAACGATAGGCATCTCCTTCGAAACGTCGGAAGTGGGGGTGTGAGTCGGTTTAGGTTT
7561	PheGlyTyrGlyAlaLysAspValArgCysHisAlaArgLysAlaValThrHisIleAsn TTTGGTTATGGGGCAAAGACGTCCTCGTTGCCATGCCAGAAAGGCCGTAAACCCACATCAAC AAACCAATACCCCGTTTCTGTGCAGGCAACGGTACGGTCTTCCGGCATTTGGGTGTAGTTG
7621	SerValTrpLysAspLeuLeuGluAspAsnValThrProIleAspThrThrIleMetAla TCCGTGTGGAAGACCTTCTGTGGAAGACAAATGTAAACCAATAGACACTACCATCATGGCT AGGCACACCTTCTGTGAAGACCTTCTGTACATTTGTGGTTATCTGTGATGGTAGTACCGA
7681	LysAsnGluValPheCysValGlnProGluLysGlyGlyArgLysProAlaArgLeuIle AAGAACGAGGTTTCTGCGTTACGCTGAGAAAGGGGGTTCGTAAAGCCAGCTCGTCTCATC TTCTTGCTCCAAAGACGCAAGTCGGAAGTCTTCCCCCAGCATTCGGTCGAGCAGAGTAG
7741	ValPheProAspLeuGlyValArgValCysGluLysMetAlaLeuTyrAspValValThr GTGTTCCCCGATCTGGGCGTGCGCGTGTGCGAAAGATGGCTTTGTACGACGTGGTTACA CACAAGGGGCTAGACCCGACGCGCACACGCTTTTCTACCGAAACATGCTGCACCAATGT
7801	LysLeuProLeuAlaValMetGlySerSerTyrGlyPheGlnTyrSerProGlyGlnArg AAGCTCCCCCTTGGCCGTGATGGGAAGCTCCTACGGATTCCAATACTCACCGAGACAGCGG TTCGAGGGGAACCGGCACTACCCCTTCGAGGATGCCTAAGGTTATGAGTGGTCCCTGTGCGCC
7861	ValGluPheLeuValGlnAlaTrpLysSerLysLysThrProMetGlyPheSerTyrAsp GTTGAATTCCCTCGTGCAAGCGTGGAAGTCCAAGAAACCCCAATGGGGTTCTCGTATGAT CAACTTAAGGAGCAGCTTCGCACCTTCAGGTTCTTTTGGGGTTACCCCAAGAGCATACTA

7921 ThrArgCysPheAspSerThrValThrGluSerAspIleArgThrGluGluAlaIleTyr
 ACCCGCTGCTTGTACTCCACAGTCACTGAGAGCGACATCCGTACGGAGGAGCAATCTAC
 TGGGCGACGAAACTGAGGTGTCACTGCTCGCTGTAGGCATGCCTCCTCCGTTAGATG

 7981 GlnCysCysAspLeuAspProGlnAlaArgValAlaIleLysSerLeuThrGluArgLeu
 CAATGTTGTGACCTCGACCCCAAGCCCGCGTGGCCATCAAGTCCCTCACCGAGAGGCTT
 GTTACAACACTGGAGCTGGGGGTTCGGGCGCACCGGTAGTTCAGGGAGTGGCTCTCCGAA

 8041 TyrValGlyGlyProLeuThrAsnSerArgGlyGluAsnCysGlyTyrArgArgCysArg
 TATGTTGGGGCCCTCTTACC AATTCAAGGGGGAGAACTGCGGCTATCGCAGGTGCCGC
 ATACAACCCCGGAGAAATGGTTAAGTTCCCCCTCTTGACGCCGATAGCGTCCACGGCG

 8101 AlaSerGlyValLeuThrThrSerCysGlyAsnThrLeuThrCysTyrIleLysAlaArg
 GCGAGCGGCGTACTGACAACTAGCTGTGGTAACACCCCTCACTTGCTACATCAAGGCCCGG
 CGCTCGCCGATGACTGTTGATCGACACCATTTGTGGGAGTGAAACGATGTAGTTCCGGGCC

 8161 AlaAlaCysArgAlaAlaGlyLeuGlnAspCysThrMetLeuValCysGlyAspAspLeu
 GCAGCCTGTCGAGCCGCGAGGGCTCCAGACTGCACCATGCTCGTGTGTGGCGACGACTTA
 CGTCGGACAGCTCGGCGTCCCGAGGTCCTGACGTGTACGAGCACACACCGCTGCTGAAT

 8221 ValValIleCysGluSerAlaGlyValGlnGluAspAlaAlaSerLeuArgAlaPheThr
 GTCGTTATCTGTGAAGCGCGGGGTCCAGGAGGACGCGGCGAGCCTGAGAGCCTTCACG
 CAGCAATAGACACTTTCGGGCCCCCAGGTCCCTCGGCCCGCTCGGACTCTCGGAAGTGC

FIG. 1V

FIG. 1W

8281 GluAlaMetThrArgTyrSerAlaProProGlyAspProProGlnProGluTyrAspLeu
 GAGGCTATGACCAGGTACTCCGCCCCCTGGGACCCCCACAAACCAGATACTG
 CTCGGATACTGGTCCATGAGCGGGGGACCCCTGGGGGTGTGGTCTTATGCTGAAC

 8341 GluLeuIleThrSerCysSerSerAsnValSerValAlaHisAspGlyAlaGlyLysArg
 GAGCTCATAACATCATGCTCCTCCTCCAACGTGTCTAGTCGCCACGACGGCGCTGGAAGAGG
 CTCGAGTATTGTAGTACGAGGAGGTGCACAGTCAGCGGGTCTGCCGCCACCTTTCTCC

 8401 ValTyrTyrLeuThrArgAspProThrThrProLeuAlaArgAlaAlaTrpGluThrAla
 GTCTACTACCTCACCCGTGACCCCTACAACCCCTCGCGAGAGCTGCGTGGAGACAGCA
 CAGATGATGGAGTGGGCACTGGGATGTTGGGGGAGCGCTCTCGACGCACCCCTCTGTCTGT

 8461 ArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMetPheAlaProThrLeuTrp
 AGACACACTCCAGTCAATTCTCTGGCTAGGCAACATAATCATGTTCGCCCCACACTGTGG
 TCTGTGTGAGGTCAGTTAAGGACCGATCCGTTGTATTAGTACAAACGGGGTGTGACACC

 8521 AlaArgMetIleLeuMetThrHisPhePheSerValLeuIleAlaArgAspGlnLeuGlu
 GCGAGGATGATACTGATGACCCATTCTTTAGCGTCCTTATAGCCAGGACCACTTGAA
 CGCTCCTACTATGACTACTGGGTAAGAAATCGCAGGAATATCGGTCCCTGGTCGAACTT

 8581 GlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIleGluProLeuAspLeuPro
 CAGGCCCTCGATTGCGAGATCTACGGGCTGCTACTCCATAGAACCACTTGATCTACCT
 GTCCGGGAGCTAACGCTCTAGATGCCCCGGACGATGAGGTATCTTGGTGAACTAGATGGA

 8641 ProIleIleGlnArgLeuHisGlyLeuSerAlaPheSerLeuHisSerTyrSerProGly
 CCAATCATTCAAAGACTCCATGGCCTCAGCGCATTTTCACTCCACAGTTACTCTCCAGGT
 GGTTAGTAAGTTTCTGAGGTACCGGAGTCCGGTAAAGTGAGGTGTCAATGAGAGGTCCA

8701 GluIleAsnArgValAlaAlaCysLeuArgLysLeuGlyValProProLeuArgAlaTrp
 GAAATTAATAGGTGGCCGTCATGCCCTCAGAAACTTGGGGTACCGCCCTTGGAGCTTGG
 CTTTAATTATCCCAACCGCGGTACGGAGTCTTTTGAACCCCATGGCGGGAACGCTCGAACC

 Gly
 8761 ArgHisArgAlaArgSerValArgAlaArgLeuLeuAlaArgGlyGlyArgAlaIle
 AGACACCGGGCCCGAGCGTCCGCGCTAGGCTTCTGGCCAGAGGAGGAGGCTGCCATA
 TCTGTGGCCCGGCGCTCGCAGGCGCATCCGAAGACCGGTCTCCTCCGTCCCGACGGTAT

 8821 CysGlyLysTyrLeuPheAsnTrpAlaValArgThrLysLeuLysLeuThrProIleAla
 TGTGGCAAGTACCTCTTCAACTGGGCGAGTAAGAAACAAAGCTCAAACTCACTCCAATAGCG
 ACACCGTTTCATGGAGAAAGTTGACCCCGTCATTCTTGTTCGAGTTTGAGTGAGGTATCGC

 8881 AlaAlaGlyGlnLeuAspLeuSerGlyTrpPheThrAlaGlyTyrSerGlyGlyAspIle
 GCCGCTGGCCAGCTGGACTTGTCCGGCTGGTTCACGGCTGGCTACAGCGGGGAGACATT
 CGCGACCGGTCGACCTGAACAGGCCGACCAAGTGCCGACCGATGTGCGCCCTCTGTAA

 (Pro)
 8941 TyrHisSerValSerHisAlaArgProArgTrpIleTrpPheCysLeuLeuLeuAla
 TATCACAGCGTGTCATGCCCGGCCCGCTGGATCTGGTTTTCCTACTCCTGCTTGCT
 ATAGTGTGCGACAGAGTACGGGCGGGCGGACCTAGACCAAAACGGATGAGGACGAACGA

FIG. 1X

FIG. 1Y

9001 AlaGlyValGlyIleTyrLeuLeuProAsnArgOP
GCAGGGGTAGGCATCTACCTCCTCCCAACCGATGAAGTTGGGGTAAACACTCCGGCCT
CGTCCCCATCCGTAGATGGAGGGGTGGCTACTTCCAAACCCCATTTGTGAGGCCCGGA

() = Heterogeneity due possibly to 5' or 3'-
terminal cloning artefact




FIG. 2